

## Higgs amplitude mode in the vicinity of a (2 + 1)-dimensional quantum critical point

A. Rançon<sup>1</sup> and N. Dupuis<sup>2</sup>

<sup>1</sup>James Franck Institute and Department of Physics, University of Chicago, Chicago, Illinois 60637, USA

<sup>2</sup>Laboratoire de Physique Théorique de la Matière Condensée, CNRS UMR 7600, Université Pierre et Marie Curie, 4 Place Jussieu, 75252 Paris Cedex 05, France

(Received 14 February 2014; revised manuscript received 16 April 2014; published 6 May 2014)

We study the “Higgs” amplitude mode in the relativistic quantum  $O(N)$  model in two space dimensions. Using the nonperturbative renormalization group we compute the  $O(N)$ -invariant scalar susceptibility in the vicinity of the zero-temperature quantum critical point. In the zero-temperature ordered phase, we find a well-defined Higgs resonance for  $N = 2$  with universal properties in agreement with quantum Monte Carlo simulations. The resonance persists at finite temperature below the Berezinskii-Kosterlitz-Thouless transition temperature. In the zero-temperature disordered phase, we find a maximum in the spectral function which is, however, not related to a putative Higgs resonance. Furthermore we show that the resonance is strongly suppressed for  $N \geq 3$ .

DOI: [10.1103/PhysRevB.89.180501](https://doi.org/10.1103/PhysRevB.89.180501)

PACS number(s): 05.30.Rt, 05.30.Jp, 67.85.-d, 75.10.-b

*Introduction.* At low temperatures many condensed-matter systems are described by a relativistic effective field theory with  $O(N)$  symmetry ( $N \geq 2$ ): quantum antiferromagnets, superconductors, Bose-Einstein condensates in optical lattices, etc. Far from criticality, collective excitations in these systems are in general well understood. In the disordered (symmetric) phase there are  $N$  gapped modes. In the ordered phase, where the  $O(N)$  symmetry is spontaneously broken, there are  $N - 1$  gapless Goldstone modes corresponding to fluctuations of the direction of the  $N$ -component quantum field, and a gapped amplitude (“Higgs”) mode [1].

The fate of the Higgs mode in low-dimensional systems near a (zero-temperature) quantum critical point (QCP) has been a subject of debate. Does the Higgs mode exist as a resonancelike feature or is it overdamped due to its coupling to Goldstone modes? In three dimensions, where the effective field theory is four-dimensional and noninteracting at the QCP, the Higgs resonance becomes sharper and sharper as the QCP is approached. This has been beautifully confirmed in the quantum antiferromagnet  $\text{TiCuCl}_3$  [2] (see also Ref. [3] for an experiment with cold atoms). In two space dimensions, the effective field theory is strongly coupled at the QCP and the existence of the Higgs resonance is not guaranteed. Furthermore the visibility of the Higgs mode strongly depends on the symmetry of the probe [4]. The longitudinal susceptibility is dominated by the Goldstone modes and diverges as  $1/\omega$  at low frequencies [5–7], thus making the observation of the Higgs resonance impossible. The  $O(N)$ -invariant scalar susceptibility (i.e., the correlation function of the square of the order parameter field) has a spectral weight which vanishes as  $\omega^3$  and is a much better candidate [4]. Quantum Monte Carlo (QMC) simulations of (2 + 1)-dimensional systems have shown that the Higgs resonance shows up in the scalar susceptibility and remains a well-defined excitation arbitrarily close to the QCP for  $N = 2$  and  $N = 3$  [8–11]. A signature of the Higgs mode has recently been observed in a two-dimensional Bose gas in an optical lattice in the vicinity of the superfluid–Mott-insulator transition [12].

In this Rapid Communication, we use a nonperturbative renormalization-group (NPRG) approach to the relativistic quantum  $O(N)$  model and compute the scalar susceptibility near the QCP. We obtain the spectral function for arbitrary val-

ues of  $N$ . For  $N = 2$ , we find a well-defined Higgs resonance in the zero-temperature ordered phase with universal properties in good agreement with QMC simulations of a related model [8,9] and the Bose-Hubbard model [10,11]. The resonance persists at finite temperature below the Berezinskii-Kosterlitz-Thouless (BKT) transition temperature; finite-temperature effects modify the spectral function only at low frequencies  $\omega \lesssim T$ . Although we also find a maximum in the spectral function in the zero-temperature disordered phase, our RG analysis shows that this maximum cannot be interpreted as a Higgs resonance as recently suggested [11]. Furthermore, we find that the resonance is strongly suppressed for  $N \geq 3$ .

*Methods.* We consider the relativistic quantum  $O(N)$  model defined by the (Euclidean) action

$$S[\boldsymbol{\varphi}, h] = \int dx \left\{ \frac{1}{2} (\nabla \boldsymbol{\varphi})^2 + \frac{1}{2c^2} (\partial_\tau \boldsymbol{\varphi})^2 + \frac{r_0}{2} \boldsymbol{\varphi}^2 + \frac{u_0}{4!} (\boldsymbol{\varphi}^2)^2 - h \boldsymbol{\varphi}^2 \right\}, \quad (1)$$

where we use the notations  $x = (\mathbf{r}, \tau)$  and  $\int dx = \int_0^\beta d\tau \int d^2r$ . The  $N$ -component real field  $\boldsymbol{\varphi}$  is periodic in the imaginary time  $\tau$ :  $\boldsymbol{\varphi}(\mathbf{r}, \tau + \beta) = \boldsymbol{\varphi}(\mathbf{r}, \tau)$  ( $\beta = 1/T$  and we set  $\hbar = k_B = 1$ ).  $r_0$  and  $u_0$  are temperature-independent coupling constants and  $c$  is the (bare) velocity of the  $\boldsymbol{\varphi}$  field. We have added a uniform time-dependent source  $h(\tau)$  which couples to  $\boldsymbol{\varphi}(x)^2$  (and will eventually be set to zero). The model is regularized by an ultraviolet cutoff  $\Lambda$  acting both on momenta and frequencies.

In two space dimensions, the phase diagram of the relativistic quantum  $O(N)$  model is well known (Fig. 1) [1]. At zero temperature (and for  $h = 0$ ), there is a quantum phase transition between a disordered phase ( $r_0 > r_{0c}$ ) and an ordered phase ( $r_0 < r_{0c}$ ) where the  $O(N)$  symmetry of the action (1) is spontaneously broken ( $u_0$  and  $c$  are considered as fixed parameters). The QCP at  $r_0 = r_{0c}$  is in the universality class of the three-dimensional classical  $O(N)$  model with a dynamical critical exponent  $z = 1$  (this value follows from Lorentz invariance at zero temperature); the phase transition is governed by the three-dimensional Wilson-Fisher fixed point. At finite temperatures, the system is always disordered,




FIG. 1. Phase diagram of the relativistic quantum  $O(N)$  model in two space dimensions ( $N \geq 2$ ). The thick line shows the zero-temperature ordered phase with long-range order (LRO), while the dashed lines are crossover lines between the renormalized classical (RC), quantum critical (QC), and quantum disordered (QD) regimes. The dotted line shows the limit of the high- $T$  region where the physics is not controlled by the QCP anymore. (For  $N = 2$ , there is a finite-temperature BKT transition line for  $r_0 \leq r_{0c}$ , which terminates at  $T = 0$  for  $r_0 = r_{0c}$ .)

in agreement with the Mermin-Wagner theorem, but it is possible to distinguish three regimes in the vicinity of the QCP depending on the temperature dependence of the correlation length  $\xi(T)$ : a renormalized classical regime where  $\xi(T) \sim (c/\rho_s)e^{2\pi\rho_s/(N-2)T}$  (with  $\rho_s$  the zero-temperature “stiffness”), a quantum critical regime where  $\xi(T) \sim c/T$ , and a quantum disordered regime where  $\xi(T) \sim \xi(0)$  [1,13]. For  $N = 2$  and  $r_0 < r_{0c}$ , there is a finite-temperature Berezinskii-Kosterlitz-Thouless (BKT) phase transition [14] and the system exhibits algebraic order at low temperatures. The BKT transition temperature line  $T_{\text{BKT}}$  terminates at the QCP  $r_0 = r_{0c}$ .

The zero-momentum scalar susceptibility is defined by

$$\begin{aligned} \chi(i\omega_n) &= \int dx e^{i\omega_n \tau} [\langle \phi(x)^2 \phi(0)^2 \rangle - \langle \phi(0)^2 \rangle^2] \\ &= \left. \frac{\delta^2 \ln Z[h]}{\delta h(-i\omega_n) \delta h(i\omega_n)} \right|_{h=0}, \end{aligned} \quad (2)$$

where  $Z[h]$  is the partition function obtained from the action (1) and  $h(i\omega_n)$  the Fourier transform of  $h(\tau)$ . The spectral function

$$\chi''(\omega) = \text{Im}[\chi(i\omega_n \rightarrow \omega + i0^+)] \quad (3)$$

is obtained by analytical continuation from Matsubara frequencies  $\omega_n = 2n\pi T$  ( $n$  integer) to real frequencies  $\omega$ . At zero temperature and in the universal regime near the QCP (scaling limit) [15],  $\chi''$  takes the form [16]

$$\chi''(\omega) = \mathcal{A}_{\pm} \Delta^{3-2/\nu} \Phi_{\pm}\left(\frac{\omega}{\Delta}\right), \quad (4)$$

where the index  $+/-$  refers to the disordered and ordered phases, respectively. In the disordered phase,  $\Delta \propto (r_0 - r_{0c})^\nu$  is the gap in the excitation spectrum (with  $\nu$  the correlation-length exponent at the QCP). In the ordered phase,  $\Delta$  is defined as the gap at the mirror point (with respect to the QCP) in the disordered phase; the ratio  $\Delta/\rho_s$  between  $\Delta$  and the stiffness  $\rho_s$  is an  $N$ -dependent universal number [17].  $\mathcal{A}_{\pm}$  is a nonuniversal cutoff-dependent constant, while  $\Phi_{\pm}(x)$  is a universal scaling function. For  $\omega \gg \Delta$ ,  $\chi''(\omega) \sim \omega^{3-2/\nu}$  is independent of  $\Delta$ . In the ordered phase, the low-energy behavior  $\omega \ll \Delta$  is entirely determined by the Goldstone modes and  $\chi''(\omega) \sim \omega^3$  [16]. In

the disordered phase, the system is gapped and  $\chi''(\omega)$  vanishes for  $\omega < 2\Delta$ . [Since  $\chi''(\omega) = -\chi''(-\omega)$  is odd, we discuss only the positive frequency part.]

To implement the NPRG approach, we add to the action (1) an infrared regulator term  $\Delta S_k[\phi]$  indexed by a momentum scale  $k$  such that fluctuations are smoothly taken into account as  $k$  is lowered from the microscopic scale  $\Lambda$  down to zero [18–20]. This allows us to introduce the scale-dependent effective action

$$\Gamma_k[\phi, h] = -\ln Z_k[\mathbf{J}, h] + \int dx \mathbf{J} \cdot \phi - \Delta S_k[\phi], \quad (5)$$

defined as a modified Legendre transform of the free energy  $-\ln Z_k[\mathbf{J}, h]$  that includes the subtraction of  $\Delta S_k[\phi]$ . Here  $\phi(x) = \delta \ln Z_k[\mathbf{J}, h]/\delta \mathbf{J}(x) = \langle \phi(x) \rangle$  is the order parameter and  $\mathbf{J}$  an external source which couples linearly to the  $\phi$  field. The variation of the effective action with  $k$  is given by Wetterich’s equation [21]

$$\partial_k \Gamma_k[\phi, h] = \frac{1}{2} \text{Tr} \{ \partial_k R_k (\Gamma_k^{(2)}[\phi, h] + R_k)^{-1} \}, \quad (6)$$

where  $\Gamma_k^{(2)}[\phi, h]$  denotes the second-order functional derivative of  $\Gamma_k[\phi, h]$  with respect to  $\phi$ . In Fourier space, the trace involves a sum over momenta and Matsubara frequencies as well as the  $O(N)$  index of the  $\phi$  field.  $R_k$  is a momentum-frequency dependent cutoff function appearing in the definition of the regulator term  $\Delta S_k[\phi]$ . The scalar susceptibility  $\chi_k(i\omega_n)$  [Eq. (2)] is obtained from the second-order functional derivative of  $\Gamma_k[\phi, h]$  with respect to the source  $h$  [22].

We solve the flow equation (6) using two main approximations. The  $h = 0$  part of the effective action  $\Gamma_k[\phi, h]$  is solved using the Blaizot-Mendez-Wschebor approximation [23–25] combined with a derivative expansion [18,19]. As for the  $h$ -dependent part, we use the following truncation:

$$\begin{aligned} \Gamma_k[\phi, h] &= \Gamma_k[\phi, 0] + \frac{1}{2} \int dy dy' h(y) H_k^{(0,2)}(y - y') h(y') \\ &\quad + \int dx dy [\rho(x) - \rho_{0,k}] H_k^{(2,1)}(x - y) h(y), \end{aligned} \quad (7)$$

where we have introduced the  $O(N)$  invariant  $\rho = \phi^2/2$  and its value  $\rho_{0,k}$  at the minimum of  $\Gamma_k[\phi, h = 0]$ . We refer to the Supplemental Material for more details on the NPRG calculation of the scalar susceptibility [22].

By numerically solving the flow equation for a given set of microscopic parameters ( $\Lambda$ ,  $u_0$ ,  $r_0$ , etc.), we obtain the scalar susceptibility  $\chi(i\omega_n) \equiv \chi_{k=0}(i\omega_n)$ . In practice, we compute  $\chi(i\omega_n)$  for typically 50 or 100 frequency points and then use a Padé approximant to deduce the spectral function  $\chi''(\omega)$  [26]. (For previous implementations of this method to compute dynamical correlation functions, see Refs. [7,27–29].)

*Higgs resonance in the ordered phase.* Figure 2 shows the scalar susceptibility  $\chi(i\omega_n)$  in the  $T = 0$  ordered phase for  $N = 2$  and various values of  $r_0$ . By plotting  $\Delta^{-\theta} [\chi(i\omega_n) - \chi(0)]$  as a function of the rescaled Matsubara frequency  $\omega_n/\Delta$ , we observe a data collapse in agreement with the expected universality. The subtraction of  $\chi(0)$  eliminates a nonsingular nonuniversal constant [8]. We expect the exponent  $\theta$  to be equal to  $3 - 2/\nu \simeq 0.02248$  [using  $\nu \simeq 0.6717$  for the three-dimensional  $O(2)$  model]; within our approximations we find  $\theta \simeq 0.1361$ . Higher-order truncations of the effective action




FIG. 2. (Color online)  $\Delta^{-\theta}[\chi(i\omega_n) - \chi(0)]$  vs  $\omega_n/\Delta$  in the  $T = 0$  ordered phase ( $N = 2$ ) for various values of  $r_0$  ( $0.001 \leq r_{0c} - r_0 \leq 0.005$  and  $\Lambda = 1$ ).

$\Gamma_k[\phi, h]$  change the value of  $\theta$  but hardly the shape of the Higgs resonance in the spectral function.

The spectral function for  $N = 2$  is shown in Fig. 3 for various values of  $r_0$  and  $T = 0$ . Again we observe a data collapse when  $\Delta^{-\theta}\chi''(\omega)$  is plotted as a function of  $\omega/\Delta$ . In the low-energy limit, the flow equation (6) is fully determined by the Goldstone mode, which leads to  $\chi''(\omega) \propto \omega^3$  as observed in Fig. 3 (see inset). For  $\omega \gg \Delta$ , we find the critical scaling  $\chi''(\omega) \sim \omega^\theta$  with  $\theta$  the exponent determined from the scaling of  $\chi(i\omega_n)$  (Fig. 2). There is a well-defined Higgs resonance whose location  $\omega = m_H$  and full width at half maximum vanish as the QCP is approached ( $r_0 \rightarrow r_{0c}$ ). The universal ratio  $m_H/\Delta \simeq 2.4$  is compatible with the QMC estimates 2.1(3) [8] and 3.3(8) [11]. Up to a multiplicative factor which depends on the nonuniversal prefactor  $\mathcal{A}_-$  (and was determined neither in the QMC simulations [8,11] nor in the present approach), the shape of the resonance, given by the universal scaling function  $\Phi_-$ , is in very good agreement with the QMC result of Ref. [8]. This gives strong support to the validity of our NPRG approach.

Figure 3 shows that the resonance persists at finite temperatures below the BKT transition temperature when  $T \ll T_{\text{BKT}} \simeq 0.42\Delta$ . At frequencies  $\omega \gg T$ , temperature has no noticeable effect: The behavior of the system is essentially quantum and the spectral function (including the Higgs resonance near  $\omega \sim m_H \gg T$ ) is well approximated by its  $T = 0$




FIG. 3. (Color online) Left: spectral function  $\Delta^{-\theta}\chi''(\omega)$  vs  $\omega/\Delta$  in the  $T = 0$  ordered phase ( $N = 2$ ) for various values of  $r_0$  ( $0.001 \leq r_{0c} - r_0 \leq 0.005$  and  $\Lambda = 1$ ). The inset shows the  $\omega^3$  dependence at small frequencies. Right: spectral function at finite temperature below the BKT transition temperature.




FIG. 4. (Color online)  $\Delta^{-\theta}\chi''(\omega)/N$  vs  $\omega/\Delta$  for various values of  $N$  in the  $T = 0$  ordered phase.

value. At frequencies  $\omega \ll T$ , the system behaves classically and the  $\omega^3$  dependence of the spectral function at  $T = 0$  is modified. In this frequency range, the numerical procedure to perform the analytic continuation becomes questionable. Nevertheless, noting that the spectral function is dominated by the Goldstone mode when  $\omega \ll \Delta$ , we can use perturbation theory to obtain

$$\chi''(\omega) \sim \omega^3 \coth\left(\frac{\omega}{2T}\right) \quad (8)$$

for  $\omega, T \ll \Delta$ . We conclude that  $\chi''(\omega) \sim T\omega^2$  at low frequencies  $\omega \ll T$ .

In Fig. 4 we show that the Higgs resonance is significantly suppressed for  $N \geq 3$  (when we vary  $N$  continuously between 2 and 3 we observe a gradual suppression). QMC simulations predict that the resonance is still marked for  $N = 3$  [8,9]. This discrepancy could be due to the limited precision of our method and more refined calculations are necessary to reach a definite conclusion regarding the precise form of  $\Phi_-(x)$ . For  $N = 1000$  we recover the exact  $N \rightarrow \infty$  result [16] showing no sign of a Higgs resonance.

*Absence of Higgs resonance in the  $T = 0$  disordered phase.* Figure 5 shows the spectral function  $\chi''(\omega)$  in the zero-temperature disordered phase for  $N = 2$ .  $\chi''(\omega)$  vanishes for  $\omega < 2\Delta$ , rises sharply above the threshold, and exhibits a maximum for  $\omega \simeq 2.5\Delta$ . Again these results are in good agreement with QMC simulations [8–11]. It has been suggested that the maximum observed in the spectral function can be interpreted as a Higgs resonance even though the




FIG. 5. (Color online)  $\Delta^{-\theta}\chi''(\omega)$  vs  $\omega/\Delta$  in the  $T = 0$  disordered phase ( $N = 2$ ) for various values of  $r_0$  ( $0.001 \leq r_{0c} - r_0 \leq 0.005$  and  $\Lambda = 1$ ).




FIG. 6. (Color online)  $\chi_k''(\omega)$  vs  $\omega/\Delta$  for various values of  $t = \ln(k/\Lambda)$  ( $T = 0$  and  $N = 2$ ). Left panel: ordered phase ( $t_J = -\ln(\xi_J \Delta) \simeq -5.2$ ). Right panel: disordered phase ( $\rho_{0,k}$  vanishes for  $t \simeq -3.9$ ).

system is disordered [10,11,30]. However, the behavior at length scales smaller than the correlation length  $\xi$  is typical of a critical system [31,32]; at no length scales does the system behave as if it were ordered. This makes the  $T = 0$  disordered phase fundamentally different from the finite-temperature phase below the BKT transition even though both phases are characterized by the absence of true long-range order.

To illustrate this point, let us discuss the spectral function  $\chi_k''(\omega)$  as a function of the renormalization group (RG) momentum scale  $k$ . In the ordered phase, the RG flow exhibits a crossover between a critical regime  $k \gg \xi_J^{-1}$  and a Goldstone regime  $k \ll \xi_J^{-1}$ , where  $\xi_J = c/\rho_s \sim c/\Delta$  is the Josephson length [17]. Figure 6 shows that the Higgs resonance is absent in the critical regime of the flow  $k \gg \xi_J^{-1}$  and quickly builds up once  $k$  reaches the Josephson scale  $\xi_J^{-1}$ . At nonzero but small temperature,  $T \ll T_{\text{BKT}}$ , the flow exhibits a

quantum-classical crossover for  $k \sim T/c \ll \xi_J^{-1}$  which modifies the low-frequency behavior of the spectral function but not the Higgs resonance.

In the disordered phase, the flow is critical as long as  $\rho_{0,k}$  (the minimum of the effective action  $\Gamma_k[\phi]$ ) remains nonzero. At the value  $k = k_c \sim \xi^{-1}$  for which  $\rho_{0,k}$  vanishes, the maximum above the gap is still not formed. It builds up for  $k < k_c$  when the RG flow is in the disordered phase ( $\rho_{0,k} = 0$ ). This definitively rules out the interpretation of the peak in the spectral function as a Higgs resonance. By the same argument we can also rule out the existence of a Higgs resonance in the finite-temperature disordered phase (quantum critical and quantum disordered regimes).

*Conclusion.* In summary, we have calculated the scalar susceptibility  $\chi(\omega)$  in the vicinity of a  $(2+1)$ -dimensional QCP. Besides the confirmation that a Higgs resonance is present for  $N = 2$  in the  $T = 0$  ordered phase, and the calculation of the universal properties of the spectral function  $\chi''(\omega)$  in agreement with quantum Monte Carlo simulations, we have shown that the resonance persists at finite temperatures below the BKT transition temperature. The spectral function then shows a  $T\omega^2$  dependence for  $\omega \ll T$ . We have also studied the possibility that a Higgs resonance exists in the zero- or finite-temperature disordered phase (characterized by a finite correlation length  $\xi$ ): Our RG analysis unambiguously reveals that the maximum observed in the spectral function cannot be interpreted as a Higgs resonance. Finally, we have shown that the resonance is strongly suppressed when  $N \geq 3$ . For  $N = 3$ , our result disagrees with Monte Carlo simulations and more refined calculations are necessary to reach a definite conclusion.

*Acknowledgments.* We thank D. Podolsky and S. Gazit for useful correspondence.

- 
- [1] S. Sachdev, *Quantum Phase Transitions*, 2nd ed. (Cambridge University Press, Cambridge, England, 2011).
- [2] C. Rüegg, B. Normand, M. Matsumoto, A. Furrer, D. F. McMorrow, K. W. Krämer, H. U. Güdel, S. N. Gvasaliya, H. Mutka, and M. Boehm, *Phys. Rev. Lett.* **100**, 205701 (2008).
- [3] U. Bissbort, S. Götze, Y. Li, J. Heinze, J. S. Krauser, M. Weinberg, C. Becker, K. Sengstock, and W. Hofstetter, *Phys. Rev. Lett.* **106**, 205303 (2011).
- [4] D. Podolsky, A. Auerbach, and D. P. Arovas, *Phys. Rev. B* **84**, 174522 (2011).
- [5] S. Sachdev, *Phys. Rev. B* **59**, 14054 (1999).
- [6] W. Zwerger, *Phys. Rev. Lett.* **92**, 027203 (2004).
- [7] N. Dupuis, *Phys. Rev. A* **80**, 043627 (2009).
- [8] S. Gazit, D. Podolsky, and A. Auerbach, *Phys. Rev. Lett.* **110**, 140401 (2013).
- [9] S. Gazit, D. Podolsky, A. Auerbach, and D. P. Arovas, *Phys. Rev. B* **88**, 235108 (2013).
- [10] L. Pollet and N. Prokof'ev, *Phys. Rev. Lett.* **109**, 010401 (2012).
- [11] K. Chen, L. Liu, Y. Deng, L. Pollet, and N. Prokof'ev, *Phys. Rev. Lett.* **110**, 170403 (2013).
- [12] M. Endres, T. Fukuhara, D. Pekker, M. Cheneau, P. Schau, C. Gross, E. Demler, S. Kuhr, and I. Bloch, *Nature (London)* **487**, 454 (2012).
- [13] S. Chakravarty, B. I. Halperin, and D. R. Nelson, *Phys. Rev. B* **39**, 2344 (1989).
- [14] V. L. Berezinskii, *Sov. Phys. JETP* **32**, 493 (1970); **34**, 610 (1971); J. M. Kosterlitz and D. J. Thouless, *J. Phys. C* **6**, 1181 (1973); **7**, 1046 (1974).
- [15] The universal regime is defined by  $|\mathbf{p}|, |\omega|/c \ll k_G$ , where  $k_G^{-1} \sim 24\pi/(cu_0)$  is the Ginzburg length.
- [16] D. Podolsky and S. Sachdev, *Phys. Rev. B* **86**, 054508 (2012).
- [17] A. Rançon, O. Kodio, N. Dupuis, and P. Lecheminant, *Phys. Rev. E* **88**, 012113 (2013).
- [18] J. Berges, N. Tetradis, and C. Wetterich, *Phys. Rep.* **363**, 223 (2002).
- [19] B. Delamotte, in *Renormalization Group and Effective Field Theory Approaches to Many-Body Systems*, edited by A. Schwenk and J. Polonyi, Lecture Notes in Physics Vol. 852 (Springer, Berlin, Heidelberg, 2012), pp. 49–132.
- [20] P. Kopietz, L. Bartosch, and F. Schütz, *Introduction to the Functional Renormalization Group* (Springer, Berlin, 2010).
- [21] C. Wetterich, *Phys. Lett. B* **301**, 90 (1993).
- [22] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevB.89.180501> for details on the NPRG calculation of the scalar susceptibility.
- [23] J.-P. Blaizot, R. Méndez-Galain, and N. Wschebor, *Phys. Lett. B* **632**, 571 (2006).

- [24] F. Benitez, J. P. Blaizot, H. Chaté, B. Delamotte, R. Méndez-Galain, and N. Wschebor, *Phys. Rev. E* **80**, 030103(R) (2009).
- [25] F. Benitez, J.-P. Blaizot, H. Chaté, B. Delamotte, R. Méndez-Galain, and N. Wschebor, *Phys. Rev. E* **85**, 026707 (2012).
- [26] H. J. Vidberg and J. W. Serene, *J. Low Temp. Phys.* **29**, 179 (1977).
- [27] N. Dupuis, *Phys. Rev. Lett.* **102**, 190401 (2009).
- [28] A. Sinner, N. Hasselmann, and P. Kopietz, *Phys. Rev. Lett.* **102**, 120601 (2009).
- [29] A. Sinner, N. Hasselmann, and P. Kopietz, *Phys. Rev. A* **82**, 063632 (2010).
- [30] Note that this suggestion is not supported by the authors of Refs. [8,9].
- [31] The system exhibits a critical behavior at length scales  $\xi_G \ll L \ll \xi$ , where  $\xi_G = k_G^{-1}$  is the Ginzburg length [15].
- [32] No resonance can form when the system is critical, in agreement with the  $\omega^{3-2/\nu}$  behavior of the spectral function.