

Modular correspondences and replicable functions (unabridged version)

J.-M. Maillard^ℒ

^ℒ LPTMC, UMR 7600 CNRS, Sorbonne Université, Tour 23, 5ème étage, case 121, 4 Place Jussieu, 75252 Paris Cedex 05, France

E-mail: maillard@lptmc.jussieu.fr

Abstract.

Landen transformation, and more generally modular correspondences, can be seen to be exact symmetries of some integrable lattice models, like the square Ising model, or the Baxter model. They are solutions of remarkable Schwarzian equations and have some compositional properties. Most of the known examples correspond, in an elliptic curves framework, to an automorphy property of pullbacked ${}_2F_1$ hypergeometric functions, associated with modular forms. It is, however, important to underline that these Schwarzian equations go beyond an elliptic curves, and hypergeometric functions framework. The question of a modular correspondence interpretation of the solutions of these “Schwarzian” equations was clearly an open question. This paper tries to shed some light on this open question. We first shed some light on the very nature of a one-parameter series solution of the Schwarzian equation. This one-parameter series is not generically a modular correspondence series, but it actually reduces to an infinite set of modular correspondence series for an infinite set of (N -th root of unity) values of the parameter. We also provide an example of two-parameter series, with a compositional property, solution of a Schwarzian equation. We finally provide simple pedagogical examples that are very similar to modular correspondence series, but are far beyond the elliptic curves framework. These last examples show that the modular correspondence-like series, or the nome-like series, are not necessarily globally bounded. The results of that paper can be seen as an incentive to study differentially algebraic series with integer coefficients, in physics or enumerative combinatorics.

PACS: 05.50.+q, 05.90.+m, 05.10.-a, 02.30.Hq, 02.30.lk, 02.30.Gp, 02.40.Xx

AMS Classification scheme numbers: 34M55, 47Exx, 32Hxx, 32Nxx, 34Lxx, 34Mxx, 14Kxx, 14H52

Key-words: Replicable functions, modular equations, modular correspondences, modular curves, modular forms, mirror maps, Schwarzian derivative, Fuchsian linear differential equations, hypergeometric functions, Square Ising model, globally nilpotent linear differential operators, Hauptmoduls, elliptic functions, Shimura curves, automorphic forms, renormalization group, Malgrange pseudo-group.

23 May 2025

1. Introduction: infinite order symmetries.

There is no need to underline the crucial role played by the concept of symmetry in physics, theoretical physics, mathematical physics. We will not consider here continuous symmetry groups (Lie groups) but rather *discrete* symmetries, not necessarily corresponding to geometrical symmetries (Coxeter groups, Weyl groups of infinite-dimensional Kac–Moody algebras), or finite groups. At first sight we do not expect any representation of these *discrete* symmetries as *linear* transformations of vector spaces (no representation theory). Examples of such *discrete symmetries*, without representation as *linear* transformations, are, for instance, *birational transformations* [1, 2], which are known to be (infinite order) discrete symmetries of integrable models [3, 4]. Such discrete symmetries can be studied, *per se*, in a discrete dynamical perspective[‡].

The simplest example of such discrete symmetries corresponds to a (univariate) transformation $x \rightarrow y(x)$ preserving some structures[†]. These structures must be *invariant*, or *covariant*, under the previous transformations $x \rightarrow y(x)$. The simplest example of “structure” is certainly just a function. Let us consider a function $\Phi(x)$, let us discard the (too simple) invariance situation, where we have a functional equation $\Phi(y(x)) = \Phi(x)$, and let us consider the following “covariance” property for a function $\Phi(x)$

$$\Phi(y(x)) = \mathcal{A}(x) \cdot \Phi(x), \quad (1)$$

where the “automorphy” cofactor $\mathcal{A}(x)$ can be described in terms of the symmetry transformation $y(x)$. Along this line the function $\Phi(x)$ can be seen as an “automorphic” function [10] with respect to the transformation $x \rightarrow y(x)$: the composition of the transformation $y(x)$ with itself, clearly yields another “covariance” or “automorphy” property

$$\Phi(y(y(x))) = \mathcal{A}(y(x)) \cdot \Phi(y(x)) = (\mathcal{A}(y(x)) \cdot \mathcal{A}(x)) \cdot \Phi(x), \quad (2)$$

and so on, for every n -th iteration of $y(x)$ with itself. From a mathematical view-point such an “automorphy property” (1) is reminiscent of the theory of *automorphic forms* [10, 11, 12, 13] (which can be generalized to Hilbert modular forms for two, or more, variables), which generalizes the theory of *modular forms* [14, 15, 16, 17, 18, 19]. In the case where $y(x)$ is not only a rational function, but a linear fractional transformation, the “covariance” property (1) can be illustrated by the Poincaré series [20, 21, 22], and other Theta-Fuchsian functions or series [13, 23, 24, 25]. From a physics view-point such an “automorphy property” (1) is reminiscent of the renormalization group theory, revisited by Wilson [26, 27], seen as a fundamental symmetry in lattice statistical mechanics or field theory. The graduate student example of exact renormalization calculation of the partition function of the one-dimensional Ising model, displayed in [28], relies on an “automorphy relation” (1), where $\Phi(x)$ is the partition function per site, and $y(x)$ corresponds to the renormalization transformation symmetry $\tanh(K) \rightarrow \tanh(K)^2$.

In the following we will not restrict the transformation symmetry $y(x)$ to be a linear fractional transformation: the function $y(x)$ is a series, analytic at $x = 0$, it

[‡] One can recall that the theory of iteration of rational functions was seen, in the pioneering work of Julia, Fatou and Ritt, as a method for investigating functional equations [5, 6, 7, 8]

[†] These structures can be linear (or non-linear) differential equations, systems of partial differential equations [9], functional equations, etc ...

can be a rational function, an algebraic function, a D-finite function, a D-D-finite function¶, a *differentially algebraic* function††, ...

To be more specific, let us give a simple, but highly pedagogical, illustration of a “covariance” property (1), which corresponds to $\Phi(x)$ being a selected ${}_2F_1$ hypergeometric function [30, 31]

$${}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], y(x)\right) = \mathcal{A}(x) \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], x\right), \quad (3)$$

where the “automorphic prefactor” $\mathcal{A}(x)$ reads

$$\mathcal{A}(x) = \lambda \cdot \left(\frac{u(x)}{u(y(x))} \cdot y'(x) \right)^{1/2}, \quad (4)$$

and where $u(x)$ is related [30, 31] to the wronskian of the order-two linear differential operator annihilating $\Phi(x)$, namely the ${}_2F_1$ hypergeometric function ${}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], x\right)$.

1.1. Modular forms, correspondences and physics.

The simplest example of a transformation $x \rightarrow y = y(x)$ occurring in the “automorphy” relation (3), or occurring as an *exact generator* of the *renormalization group* of the square Ising model, or even of the Baxter model [32], corresponds to the *Landen transformation* [32, 33]

$$k \longrightarrow k_L = \frac{2\sqrt{k}}{1+k}, \quad (5)$$

or to its compositional inverse, the inverse Landen transformation. As it should, the *critical point* of the square Ising model (resp. Baxter model) is a *fixed point* [32] of the Landen transformation: $k = 1$.

Let us introduce the j -invariant‡ of an elliptic curve of modulus k , and its transform by the Landen transformation (5)

$$j(k) = 256 \cdot \frac{(1 - k^2 + k^4)^3}{k^4 \cdot (1 - k^2)^2}, \quad j(k_L) = 16 \cdot \frac{(1 + 14k^2 + k^4)^3}{(1 - k^2)^4 \cdot k^2}, \quad (6)$$

and let us also introduce the two corresponding *Hauptmoduls* [32]:

$$x = \frac{1728}{j(k)}, \quad y = \frac{1728}{j(k_L)}. \quad (7)$$

These two Hauptmoduls (7) are related by the *modular equation* [35, 36, 37, 38, 39, 40]:

$$\begin{aligned} 1953125 x^3 y^3 - 187500 x^2 y^2 \cdot (x + y) + 375 xy \cdot (16 x^2 - 4027 xy + 16 y^2) \\ - 64 (x + y) \cdot (x^2 + 1487 xy + y^2) + 110592 xy = 0. \end{aligned} \quad (8)$$

The algebraic function $y = y(x)$, defined from the modular curve (8), is a *multivalued function*, but we can single out the series expansion††:

$$y = \frac{1}{1728} \cdot x^2 + \frac{31}{62208} \cdot x^3 + \frac{1337}{3359232} \cdot x^4 + \frac{349115}{1088391168} \cdot x^5$$

¶ A D-D-finite function is a function solution of a linear differentiable operator with D -finite function coefficients [29].

†† A differentially algebraic function is a function solution of a non-linear differential equation of the form $P(x, y, y', y'', \dots y^{(n)}) = 0$, where P is a polynomial.

‡ The j -invariant [32, 34] (see also Klein’s modular invariant) regarded as a function of a complex variable τ (the ratio of periods), is a modular function of weight zero for $SL(2, \mathbb{Z})$.

†† This series (9) has a radius of convergence 1, even if the discriminant of the modular equation (8) which vanishes at $x = 1$, vanishes for values inside the unit radius of convergence, for instance at $x = -64/125$.

$$+ \frac{20662501}{78364164096} \cdot x^6 + \frac{1870139801}{8463329722368} \cdot x^7 + \dots \quad (9)$$

The transformation $x \rightarrow y(x) = y$, where y is given by the modular equation (8), is thus an *algebraic* transformation, corresponding to the Landen transformation (5), or to the inverse Landen transformation. The emergence of a *modular form* [41, 42, 43] corresponds to the remarkable automorphy identity (3) on the *same* hypergeometric function, but where the pullback x is changed $x \rightarrow y(x) = y$, according to the modular equation (8).

Let us consider another important modular equation. The modular equation of order three corresponding to $\tau \rightarrow 3 \cdot \tau$, or $\tau \rightarrow \tau/3$, reads¶:

$$k^4 + 12 k^3 \lambda + 6 k^2 \lambda^2 + 12 k \lambda^3 + \lambda^4 - 16 k^3 \lambda^3 - 16 k \lambda = 0. \quad (10)$$

Recalling that

$$x = \frac{27}{4} \cdot \frac{k^4 \cdot (1 - k^2)^2}{(k^4 - k^2 + 1)^3} = \frac{1728}{j(k)}, \quad y = \frac{27}{4} \cdot \frac{\lambda^4 \cdot (1 - \lambda^2)^2}{(\lambda^4 - \lambda^2 + 1)^3} = \frac{1728}{j(\lambda)}, \quad (11)$$

gives a modular equation $P(x, y) = 0$, yielding the series expansion:

$$y = \frac{x^3}{2985984} + \frac{31 x^4}{71663616} + \frac{36221 x^5}{82556485632} + \frac{29537101 x^6}{71328803586048} + \dots \quad (12)$$

Note that these two series (9) and (12) *commute*. An alternative rational parametrization of this last modular equation $P(x, y) = 0$ can be found in [19]:

$$x = \frac{1728 t}{(t + 27) \cdot (t + 3)^3} \quad \text{and:} \quad y = \frac{1728 t^3}{(t + 27) \cdot (t + 243)^3}. \quad (13)$$

Again we have an automorphy relation (3) where $y(x)$ is given by (12) with an algebraic “automorphic prefactor” $\mathcal{A}(x)$.

1.2. Schwarzian condition

More generally, the Gauss hypergeometric function ${}_2F_1([\alpha, \beta], [\gamma], x)$ is solution of the second order linear differential operator†:

$$\begin{aligned} \Omega &= D_x^2 + A(x) \cdot D_x + B(x), & \text{where:} & \quad (14) \\ A(x) &= \frac{(\alpha + \beta + 1) \cdot x - \gamma}{x \cdot (x - 1)} = \frac{u'(x)}{u(x)}, & B(x) &= \frac{\alpha \beta}{x \cdot (x - 1)}. \end{aligned}$$

An automorphy relation, like (3) but on ${}_2F_1([\alpha, \beta], [\gamma], x)$, amounts to saying that the second order linear differential operator (14), pullbacked by $x \rightarrow y(x)$, reduces to the conjugate of the linear differential operator (14). Let us assume that the pullback $y(x)$ is an algebraic series like in (9) and (12). A straightforward calculation [31] allows to find the algebraic cofactor $\mathcal{A}(x)$ in terms of the algebraic function pullback $y(x)$:

$$\mathcal{A}(x) = \lambda \cdot \left(\frac{u(x)}{u(y(x))} \cdot y'(x) \right)^{-1/2}. \quad (15)$$

Expression (15) for $\mathcal{A}(x)$ is such that the two order-two linear differential operators (of a similar form as (14)) have the same D_x coefficient. The identification of these two

¶ Legendre already knew (1824) this order three modular equation in the form $(k\lambda)^{1/2} + (k'\lambda')^{1/2} = 1$, where k and k' , and λ , λ' are pairs of complementary moduli $k^2 + k'^2 = 1$, $\lambda^2 + \lambda'^2 = 1$, and Jacobi derived that modular equation [45, 46].

† Note that $A(x)$ is the log-derivative of $u(x) = x^\gamma \cdot (1 - x)^{\alpha + \beta + 1 - \gamma}$.

linear differential operators thus corresponds (beyond (15)) to *just one (non-linear) condition* that can be rewritten (after some algebra ...) in the following *Schwarzian* form:

$$W(x) - W(y(x)) \cdot y'(x)^2 + \{y(x), x\} = 0, \quad (16)$$

where

$$W(x) = A'(x) + \frac{A(x)^2}{2} - 2 \cdot B(x), \quad (17)$$

and where $\{y(x), x\}$ denotes the *Schwarzian derivative* [47]:

$$\{y(x), x\} = \frac{y'''(x)}{y'(x)} - \frac{3}{2} \cdot \left(\frac{y''(x)}{y'(x)} \right)^2 = \frac{d}{dx} \left(\frac{y''(x)}{y'(x)} \right) - \frac{1}{2} \cdot \left(\frac{y''(x)}{y'(x)} \right)^2. \quad (18)$$

For ${}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], x\right)$, the “automorphy” condition (3) yields the Schwarzian condition (16) with:

$$W(x) = -\frac{32x^2 - 41x + 36}{72 \cdot x^2 \cdot (x-1)^2}. \quad (19)$$

The algebraic series (9) and (12), associated with different modular equations (like (8)), are both solutions of the *same* Schwarzian condition (16) with $W(x)$ given by (19). These two *modular correspondences* series (9) and (12), associated with *modular curves*, are thus *algebraic series*. Consequently, the prefactor (15) is an *algebraic function*.

1.3. One-parameter solution series of the Schwarzian condition (16)

Trying to generalize the modular equation (8), and its associated algebraic series (9), let us try to find the series of the form $a \cdot x^2 + \dots$, solutions of the Schwarzian equation (16) with $W(x)$ given by (19). It is straightforward to find that such series is, in fact, the following *one-parameter* series:

$$\begin{aligned} y_2 = & a \cdot x^2 + \frac{31 \cdot ax^3}{36} - \frac{a \cdot (5952a - 9511)}{13824} \cdot x^4 - \frac{a \cdot (14945472a - 11180329)}{20155392} \cdot x^5 \\ & + \frac{a \cdot (88746430464a^2 - 677409785856a + 338926406215)}{743008370688} \cdot x^6 + \dots \end{aligned} \quad (20)$$

which actually reduces to (9) for $a = 1/1728$. Similarly, one also finds a one-parameter family of solution-series of the Schwarzian condition (16) of the form $b \cdot x^3 + \dots$, namely

$$\begin{aligned} y_3 = & b \cdot x^3 + \frac{31b}{24} \cdot x^4 + \frac{36221b}{27648} \cdot x^5 - \frac{b \cdot (23141376b - 66458485)}{53747712} \cdot x^6 \\ & - \frac{b \cdot (183649959936b - 187769367601)}{165112971264} \cdot x^7 + \dots \end{aligned} \quad (21)$$

which reduces to (12) for $b = 1/2985984 = 1/1728^2$.

Remark 1.1: Generically the two series (20) and (21) are differentially *algebraic series* (being solution of a Schwarzian condition (16), with $W(x)$ given by (19)). For *selected* values of the parameter, like $a = 1/1728$ and $b = 1/1728^2$, these series become *algebraic series* (*correspondences* associated with *modular curves*). Are there other selected values of the parameters for which the series becomes an algebraic series? Are there selected values of the parameters for which the series become (non

algebraic) D -finite series ? Are there selected values of the parameters for which the series become D - D -finite[†] series [48, 49] ?

1.4. The nome and mirror maps

Let us recall the concept of *mirror map* [41, 42, 50, 51, 52, 53, 54] relating the reciprocal of the j -function and the nome, with the well-known series with *integer* coefficients[‡] :

$$\begin{aligned}\tilde{X}(q) = & q - 744q^2 + 356652q^3 - 140361152q^4 + 49336682190q^5 \\ & - 16114625669088q^6 + 4999042477430456q^7 + \dots\end{aligned}\quad (22)$$

and the *nome* which is its compositional inverse:

$$\begin{aligned}\tilde{Q}(x) = & x + 744x^2 + 750420x^3 + 872769632x^4 + 1102652742882x^5 \\ & + 1470561136292880x^6 + 2037518752496883080x^7 + \dots\end{aligned}\quad (23)$$

The series (22) corresponds to x being the reciprocal of the j -function: $1/j$. As a consequence of the (modular form) hypergeometric identities (3) (see (7)), we need x to be identified with the *Hauptmodul* $1728/j$.

The series $X(q) = 1728 \cdot \tilde{X}(q)$ (with $\tilde{X}(q)$ given by (22)) is solution of the Schwarzian equation:

$$\{X(q), q\} - \frac{1}{2q^2} - W\left(X(q)\right) \cdot \left(\frac{dX(q)}{dq}\right)^2 = 0. \quad (24)$$

The series $Q(x) = \tilde{Q}(x/1728)$ (with $\tilde{Q}(x)$ given by (23)) is solution of the Schwarzian equation:

$$\{Q(x), x\} + \frac{1}{2 \cdot Q(x)^2} \cdot \left(\frac{dQ(x)}{dx}\right)^2 + W(x) = 0. \quad (25)$$

The two mirror map series (22), (23) thus correspond to *differentially algebraic* [55, 56] series: they are solutions of simple (non-linear) Schwarzian equations like in (16).

The two one-parameter series (20) and (21) correspond respectively to:

$$X\left(a \cdot Q(x)^2\right) \quad \text{and:} \quad X\left(b \cdot Q(x)^3\right). \quad (26)$$

More generally, all the series

$$y_n(a, x) = X\left(a \cdot Q(x)^n\right), \quad (27)$$

are solutions of the Schwarzian condition (16). For the selected values $a = 1/1728^{n-1}$ these series (27) turn out to be *algebraic series*: they are series actually associated with *correspondences*, *modular curves*. The composition of two such series is also solution of the Schwarzian condition (16). One easily finds that

$$\begin{aligned}y_n\left(a, y_m(b, x)\right) &= y_{mn}(a \cdot b^n, x) = a \cdot b^n \cdot x^{mn} + \dots \\ y_m\left(b, y_n(a, x)\right) &= y_{mn}(b \cdot a^m, x) = b \cdot a^m \cdot x^{mn} + \dots\end{aligned}\quad (28)$$

Generically the two series y_n and y_m do not commute.

[†] D -finite functions are solutions of linear differential operators with polynomial coefficients, D - D -finite series are solutions of linear differential operators with D -finite function coefficients, etc ...

[‡] In Maple the series (22) can be obtained substituting $L = \text{EllipticModulus}(q^{1/2})^2$, in $1/j = L^2 \cdot (L-1)^2/(L^2 - L + 1)^3/256$. See <https://oeis.org/A066395> for the series (22) and <https://oeis.org/A091406> for the series (23).

Remark 1.2: Do note that these two series *do commute* for the selected values $a = 1/1728^{n-1}$ and $b = 1/1728^{m-1}$, of the parameters a and b , actually associated with modular correspondences (algebraic series). In that case, one has the identity:

$$a \cdot b^n = b \cdot a^m = \frac{1}{1728^{n-1}} \cdot \left(\frac{1}{1728^{m-1}} \right)^n = \frac{1}{1728^{m \cdot n - 1}}. \quad (29)$$

Also note that if one assumes that the parameters a (resp. b) are of the form ρ^{n-1} (resp. ρ^{m-1}) with ρ different from $1/1728$ or 1 , the series $y_n(a, x)$ and $y_m(b, x)$ *still commute*[¶], even if they are *not* algebraic series but *only differentially algebraic series*. The compositional identities (29) are inherited from the fact that the composition of two algebraic series is an algebraic series, and that the *composition of two solutions of the Schwarzian condition* (16) *must† also be a solution of the Schwarzian condition* (16). Such properties are reminiscent of the concept of *replicable* functions [57, 58, 59, 60, 61, 62, 63, 64].

This set of solution series (27) of the Schwarzian condition (16), can also be obtained by the composition of algebraic series associated with modular *correspondences* (which have no parameter, see (9), (12), ...), together with the following one-parameter series $X(e \cdot Q(x))$ *also solution* of the Schwarzian condition (16). This series reads:

$$y(e, x) = e \cdot x + e \cdot (e - 1) \cdot S_e(x), \quad \text{where:} \quad (30)$$

$$S_e(x) = -\frac{31}{72} \cdot x^2 + \frac{(9907e - 20845)}{82944} \cdot x^3 - \frac{(4386286e^2 - 20490191e + 27274051)}{161243136} \cdot x^4 + \dots \quad (31)$$

Remark 1.3: It is straightforward to see that the series (30) is an order- N transformation when the parameter e is a N -th root of unity: $e^N = 1$. These N -th root of unity are, thus, clearly *selected values* of the parameters. Are all these N -th root of unity series algebraic series, or just D-finite series, or simply differentially algebraic series ?

1.5. Multivalued functions and reversibility

The Landen algebraic transformation (5) amounts to multiplying (*or dividing* because of the modular group symmetry $\tau \leftrightarrow 1/\tau$) the ratio τ of the two periods of the elliptic curves: $\tau \longleftrightarrow 2\tau$. The other (isogeny) transformations^{††} correspond to $\tau \leftrightarrow N \cdot \tau$, for various integers N .

We, thus, see that a modular equation, like (8), yields *multivalued* functions corresponding to the different series solutions of the modular equation (for instance (9) and its compositional inverse). More generally, for $\tau \leftrightarrow N \cdot \tau$, we will have series like $1/1728^{N-1} \cdot x^N + \dots$ and also (their compositional inverse Puiseux series) $1728^{(N-1)/N} \cdot x^{1/N} + \dots$.

In the textbooks the renormalization group is often presented as a semi-direct group[‡]. In fact the renormalization group generators have no reason to be such

[¶] In terms of the nome, this amounts to noticing that transformations $q \rightarrow \alpha^{n-1} \cdot q^n$ and $q \rightarrow \alpha^{m-1} \cdot q^m$ commute.

[†] This is also a clear consequence of the automorphy property (1).

^{††} See for instance (2.18) in [34].

[‡] In most of the graduate text book on renormalization group, the critical fixed point is an attractive fixed point. There is an “arrow of time”. The renormalization group is seen as an irreversible process.

irreversible transformations. They are, at first sight, *reversible transformations*. The modular equation (8) has a $x \leftrightarrow y$ symmetric polynomial, corresponding to the Landen transformation, *as well as its compositional inverse*, the inverse Landen transformation. These two transformations are both *exact generators* of the renormalization group of the square Ising model, or of the Baxter model [32]. With this exact renormalization group representation we see that the modular equation restores, *as a consequence of its $x \leftrightarrow y$ symmetry*, the *reversible character of the renormalization group*, the price to pay being that the function $y(x)$ is actually *multivalued*.

The Schwarzian condition (16) encapsulates [30, 31] an *infinite number of modular correspondences* associated with their modular curves and modular forms [14, 15, 16]. In these cases the automorphy relation (3) corresponds to *algebraic function* prefactors $\mathcal{A}(x)$. However, for series with one-parameter, like (20) and (21), which are *generically* differentially algebraic, we still have an “automorphy” relation (3), but with differentially algebraic “automorphy” prefactors $\mathcal{A}(x)$. We cannot expect a modular equation, but is there a way to still see such transformations (20) and (21), as “correspondences” with some “appropriate” generalization of the concept of correspondences ?

1.6. Correspondences, Schwarzian conditions and replicable functions

The Schwarzian condition (16) coincides exactly with one of the conditions G. Casale obtained [65, 66, 67, 68, 69, 70, 71] in a classification of Malgrange’s \mathcal{D} -envelopes and \mathcal{D} -groupoids [72] on \mathbb{P}_1 . Denoting $y'(x)$, $y''(x)$ and $y'''(x)$ the first, second and third derivative of $y(x)$ with respect to x , these conditions[‡] read respectively¶

$$\mu(y) \cdot y'(x) - \mu(x) + \frac{y''(x)}{y'(x)} = 0, \quad (32)$$

$$\nu(y) \cdot y''(x)^2 - \nu(x) + \frac{y'''(x)}{y'(x)} - \frac{3}{2} \cdot \left(\frac{y''(x)}{y'(x)} \right)^2 = 0, \quad (33)$$

together with $\gamma(y) \cdot y'(x)^n - \gamma(x) = 0$ and $h(y) = h(x)$, corresponding respectively to rank two, rank three, together with rank one and rank nul groupoids, where $\nu(x)$, $\mu(x)$, $\gamma(x)$ are *meromorphic* functions ($h(x)$ is holomorph).

The previous examples of Schwarzian condition (16) correspond to *elliptic curves* (modular curves, modular forms and modular correspondences), through pullbacked ${}_2F_1$ hypergeometric functions [19]. In subsection 3.2 of [30] we have seen that the Schwarzian condition (16) can actually occur with *Heun functions which cannot be reduced to pullbacked ${}_2F_1$ hypergeometric functions*^{††}, and which *do not* correspond to globally bounded [43, 44] series. Similarly, we have seen Schwarzian conditions (16) corresponding to (non globally bounded) pullbacked ${}_2F_1$ hypergeometric functions, associated with *Shimura curves* [75, 76]. The Malgrange-Casale approach for Schwarzian conditions (33) suggests that one should be able to find examples of such Schwarzian conditions *far beyond modular curves, or even Shimura curves* (and their associated modular forms [14, 15, 16] and automorphic forms [10]). If such generalizations exist, are they also associated with *one-parameter* series ? How

[‡] Casale’s condition (32) is *exactly the same condition* as the one we already found in [73], and this is not a coincidence.

¶ More generally see the concept of differential algebraic invariant of isogenies in [74].

^{††} See for instance the two Heun functions given by (164) in [77].

to describe them ? Can they necessarily be seen, eventually, as generalization of correspondences ?

In the next section we will first revisit the previous “classical” modular correspondence results *with a different normalization* of the pullback (see (34) below) which makes the occurrence of series with *integer* coefficients crystal clear. Revisiting these calculations with a key role played by a function $F(x)$ defined below by (170), we will be able to find some new partial differential equations (see (197), or (198) below), in the parameter of the series[†]. These new equations will help us to find many examples of replicable-like [58, 59, 60, 61, 62] functions *far beyond modular curves or Shimura curves* [75, 76].

2. Recalls

Some part of this section will be reminiscent of the results explained in [30], with the difference that we have *another normalization* of the pullback, corresponding to change $x \rightarrow 1728x$, the “automorphy” relation (3) thus becoming

$${}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y\right) = \mathcal{A}(x) \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right), \quad (34)$$

where[¶]:

$$\mathcal{A}(x) = \lambda \cdot \left(\frac{u(x)}{u(y(x))} \cdot y'(x) \right)^{1/2}. \quad (35)$$

As a consequence the (pullback) algebraic series $y = y(x)$, corresponding to isogenies like (9), (12), ... are normalized as $x \rightarrow x^N + \dots$, and are series with *integer* coefficients.

In our case, taking into account the exact expression of the wronskian, one has $u(x) = x \cdot (1 - 1728x)^{1/2}$, and, thus, we get:

$$\mathcal{A}(x) = \lambda \cdot \left(\frac{x \cdot (1 - 1728x)^{1/2}}{y \cdot (1 - 1728y)^{1/2}} \cdot y'(x) \right)^{1/2}. \quad (36)$$

Taking the square of (34) we can thus rewrite the “automorphic” relation (34) as

$$\begin{aligned} \lambda \cdot y \cdot (1 - 1728 \cdot y)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y\right)^2 \\ = x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \cdot \frac{dy}{dx}, \end{aligned} \quad (37)$$

which is, in fact, nothing but

$$\lambda \cdot \frac{dx}{F(x)} = \frac{dy}{F(y)}. \quad (38)$$

where $F(x)$ reads:

$$\begin{aligned} F(x) &= x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \\ &= x - 744x^2 - 393768x^3 - 3574444672x^4 - 394896727080x^5 + \dots \end{aligned} \quad (39)$$

The elimination of the “automorphic” cofactor $\mathcal{A}(x)$ gives the Schwarzian equation on $y(x)$

$$W(x) - W(y(x)) \cdot y'(x)^2 + \{y(x), x\} = 0, \quad (40)$$

[†] See also (229) below for more parameters.

[¶] Note a typo in (92) in [30]. the exponent $-1/2$ in (92) must be changed into $1/2$.

where now

$$W(x) = -\frac{1}{2} \cdot \frac{1 - 1968x + 2654208x^2}{x^2 \cdot (1 - 1728x)^2}, \quad (41)$$

namely:

$$\begin{aligned} & -\frac{1}{2} \cdot \frac{1 - 1968x + 2654208x^2}{x^2 \cdot (1 - 1728x)^2} \\ & + \frac{1}{2} \cdot \frac{1 - 1968y(x) + 2654208y(x)^2}{y(x)^2 \cdot (1 - 1728y(x))^2} \cdot y'(x)^2 + \{y(x), x\} = 0. \end{aligned} \quad (42)$$

3. Modular equation, modular correspondence

3.1. $q \longrightarrow q^2$

Let us consider the *modular equation*[†]:

$$\begin{aligned} \Gamma_2(x, y) = & \ x y - (x + y) \cdot (x^2 + 1487xy + y^2) \\ & + 10125 \cdot xy \cdot (16x^2 - 4027xy + 16y^2) \\ & - 8748000000 \cdot x^2 y^2 \cdot (x + y) + 15746400000000 \cdot x^3 y^3 = 0, \end{aligned} \quad (43)$$

which has the following rational parametrization [19]:

$$x = \frac{t}{(t+16)^3} \quad \text{and:} \quad y = \frac{t^2}{(t+256)^3}. \quad (44)$$

It has the following *algebraic series* solutions with *integer* coefficients

$$\begin{aligned} y_2 = & \ x^2 + 1488x^3 + 2053632x^4 + 2859950080x^5 + 4062412996608x^6 \\ & + 5882951135920128x^7 + 8664340079503736832x^8 + \dots \end{aligned} \quad (45)$$

and

$$\begin{aligned} y_{1/2} = & \ \omega \cdot x^{1/2} - 744 \cdot x^{2/2} + 357024 \cdot \omega \cdot x^{3/2} - 140914688 \cdot x^{4/2} \\ & + 49735011840 \cdot \omega \cdot x^{5/2} - 16324041375744 \cdot x^{6/2} + \dots \end{aligned} \quad (46)$$

where $\omega^2 = 1$ (i.e. $\omega = \pm 1$). These two algebraic series can be written respectively:

$$\tilde{X}(\tilde{Q}(x)^2) \quad \text{and:} \quad \tilde{X}(\omega \cdot \tilde{Q}(x)^{1/2}). \quad (47)$$

They amount, respectively, to changing the nome as follows: $q \longrightarrow q^2$, *together with its compositional inverse* $q \longrightarrow \omega \cdot q^{1/2}$, where $\omega^2 = 1$. These two series, (45) and (46), are actually solutions of the Schwarzian equation (40) with $W(x)$, now, given by (41). Note that we have the following relation:

$$\begin{aligned} & 2 \cdot y_2 \cdot (1 - 1728 \cdot y_2)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y_2\right)^2 \\ & = x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \cdot \frac{dy_2}{dx}. \end{aligned} \quad (48)$$

We have a similar relation for $y_{1/2}$. Relation (48), and the corresponding one for $y_{1/2}$, are nothing but:

$$2 \cdot \frac{dx}{F(x)} = \frac{dy_2}{F(y_2)} \quad \text{and:} \quad \frac{1}{2} \cdot \frac{dx}{F(x)} = \frac{dy_{1/2}}{F(y_{1/2})}. \quad (49)$$

[†] Which is nothing but (8) with the change of variables $x \rightarrow x/1728$, $y \rightarrow y/1728$.

3.2. Linear ODE for $q \longrightarrow q^2$

The previous *algebraic series* (45), (46) are solutions of an order-three linear differential operator $M_3 = M_1 \oplus M_2$ which is the *direct sum* (LCLM) of an order-two linear differential operator M_2 , and an order-one linear differential operator M_1 with the following rational function solution:

$$\begin{aligned} S_1^{(1)} &= \frac{x}{496} \cdot \frac{(496 + 13591125x + 2916000000x^2)}{(1 - 54000x)^3} \\ &= x + \frac{93943125}{496}x^2 + \frac{680168390625}{31}x^3 + \frac{63705259687500000}{31}x^4 + \dots \end{aligned} \quad (50)$$

Let us introduce the two formal solution series of order-two linear differential operator M_2 , namely:

$$\begin{aligned} S_1^{(2)} &= x + \frac{46971563}{248}x^2 + \frac{680168390718}{31}x^3 + \frac{63705259687628352}{31}x^4 + \dots \\ S_2^{(2)} &= x^{1/2} \cdot \left(1 + 357024x + 49735011840x^2 + 5091284519436288x^3 \right. \\ &\quad \left. + 445924637193878765568x^4 + \dots \right) \end{aligned}$$

One has the following relation

$$y_2 = 496 \cdot (S_1^{(2)} - S_1^{(1)}), \quad (51)$$

$$y_{1/2}(\omega, x) = \omega \cdot S_2^{(2)} - 248 \cdot (2S_1^{(1)} + S_1^{(2)}), \quad (52)$$

where $\omega^2 = 1$.

3.3. $q \longrightarrow q^3$

Let us consider the *modular equation*:

$$\begin{aligned} &1855425871872000000000 \cdot x^3 y^3 \cdot (y + x) \\ &+ 16777216000000 \cdot y^2 x^2 \cdot (27x^2 - 45946xy + 27y^2) \\ &+ 36864000 \cdot xy \cdot (y + x) \cdot (x^2 + 241433xy + y^2) \\ &+ (x^4 - 1069956x^3y + 2587918086x^2y^2 - 1069956xy^3 + y^4) \\ &+ 2232 \cdot xy \cdot (y + x) - xy = 0, \end{aligned} \quad (53)$$

which has the following rational parametrization [19]:

$$x = \frac{t}{(t+27) \cdot (t+3)^3} \quad \text{and:} \quad y = \frac{t^3}{(t+27) \cdot (t+243)^3}. \quad (54)$$

This *modular equation* (53) has the following *algebraic series* solutions

$$\begin{aligned} y_3 &= x^3 + 2232x^4 + 3911868x^5 + 6380013816x^6 + 10139542529238x^7 \\ &+ 15969813236020944x^8 + 25104342383076998772x^9 + \dots \end{aligned} \quad (55)$$

and its compositional inverse

$$\begin{aligned} y_{1/3}(\omega, x) &= \omega \cdot x^{1/3} - 744 \cdot \omega^2 \cdot x^{2/3} + 356652 \cdot x^{3/3} - 140360904 \cdot \omega \cdot x^{4/3} \\ &+ 49336313166 \cdot \omega^2 x^{5/3} - 16114360320000 \cdot x^{6/3} + \dots \end{aligned} \quad (56)$$

where $\omega^3 = 1$. The radius of convergence of the series (55) is $R = 1/1728$, corresponding to the vanishing of the discriminant of the modular equation (53). These two series can be written respectively

$$\tilde{X}(\tilde{Q}(x)^3) \quad \text{and:} \quad \tilde{X}(\omega \cdot \tilde{Q}(x)^{1/3}), \quad (57)$$

where $\omega^3 = 1$. They amount, respectively, to changing the nome as follows: $q \rightarrow q^3$, together with its compositional inverse $q \rightarrow \omega \cdot q^{1/3}$ where $\omega^3 = 1$. These two algebraic series, (55) and (56), are actually solutions of the Schwarzian equation (40), with $W(x)$ given by (41). Note that we have the following relation:

$$\begin{aligned} 3 \cdot y_3 \cdot (1 - 1728 \cdot y_3)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y_3\right)^2 \\ = x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \cdot \frac{dy_3}{dx}. \end{aligned} \quad (58)$$

We have a similar relation for $y_{1/3}$. Relation (58), and the corresponding one for $y_{1/3}$, are nothing but:

$$3 \cdot \frac{dx}{F(x)} = \frac{dy_3}{F(y_3)} \quad \text{and:} \quad \frac{1}{3} \cdot \frac{dx}{F(x)} = \frac{dy_{1/3}}{F(y_{1/3})}. \quad (59)$$

3.4. Linear ODE for $q \rightarrow q^3$

The previous algebraic series (55), (56) are solutions of an order-four linear differential operator $M_3 = M_1 \oplus M_3$, which is the direct sum (LCLM) of an order-three linear differential operator M_3 , and an order-one linear differential operator M_1 with the rational function solution

$$\begin{aligned} S_1^{(1)} &= \frac{x}{267489} \cdot \frac{p_3(x)}{(1 + 12288000 x)^3} \\ &= x - \frac{447621120000}{9907} x^2 + \frac{324554085892096000000}{267489} x^3 + \dots \end{aligned} \quad (60)$$

where:

$$\begin{aligned} p_3(x) &= 267489 - 2225055744000 x + 192711491584000000 x^2 \\ &\quad - 463856467968000000000 x^3. \end{aligned} \quad (61)$$

The solutions of the order-three linear differential operator M_3 read:

$$\begin{aligned} S_1^{(2)} &= x - \frac{447621120000}{9907} x^2 + \frac{10818469529736533333}{89163} x^3 + \dots, \\ S_2^{(2)} &= x^{2/3} \cdot \left(1 - \frac{8222718861}{124} x + \frac{62192008621897866}{31} x^2 \right. \\ &\quad \left. - \frac{2837950236255383813660913}{62} x^3 + \dots \right), \end{aligned} \quad (62)$$

$$\begin{aligned} S_3^{(2)} &= x^{1/3} \cdot \left(1 - 140360904 x + 4998903239356308 x^2 \right. \\ &\quad \left. - 122558022956400494032656 x^3 + \dots \right). \end{aligned} \quad (63)$$

The solutions (55) and (56) of the modular equation (53) can be expressed in terms of the solutions of the previous linear differential operators M_1 and M_3

$$y_3 = 267489 \cdot (S_1^{(1)} - S_1^{(2)}). \quad (64)$$

and:

$$y_{1/3}(\omega, x) = \omega \cdot S_3^{(2)} - 744 \cdot \omega^2 \cdot S_2^{(2)} + 89163 \cdot (3 \cdot S_1^{(1)} + S_1^{(2)}). \quad (65)$$

3.5. $q \longrightarrow q^5$

We are not going to give explicitly the modular equation corresponding to $q \longrightarrow q^5$ because it starts becoming a bit too large. Let us just say that it can (easily) be obtained by the elimination of t in its rational parameterization [19]:

$$x = \frac{t}{(t^2 + 10t + 5)^3} \quad \text{and:} \quad y = \frac{t^5}{(t^2 + 250t + 3125)^3}. \quad (66)$$

This modular curve $\Gamma_5(x, y) = \Gamma_5(y, x) = 0$, has the following *algebraic series* solutions

$$y_5 = x^5 + 3720x^6 + 9287460x^7 + 19648405600x^8 + 38124922672650x^9 \\ + 70330386411705000x^{10} + 125698841122545005000x^{11} + \dots \quad (67)$$

and

$$y_{1/5} = \omega \cdot x^{1/5} - 744 \cdot \omega^2 \cdot x^{2/5} + 356652 \omega^3 \cdot x^{3/5} - 140361152 \cdot \omega^4 \cdot x^{4/5} \\ + 49336682190 \cdot x^{5/5} - \frac{80573128344696}{5} \cdot \omega \cdot x^{6/5} + \dots \quad (68)$$

where $\omega^5 = 1$. The series (67) and (68) are (algebraic) solutions of an order-six linear differential operator $L_6 = L_1 \oplus L_5$, which is the direct sum of an order-one linear differential operator with a rational function solution ($p_5(x)$ is a polynomial with integer coefficients)

$$r(x) = \frac{x}{41113901825} \cdot \frac{p_5(x)}{1 + 654403829760x + 5209253090426880x^2} \quad (69) \\ = x - \frac{4085556703324323840000}{1644556073} x^2 + \dots$$

and an irreducible order-five linear differential operator operator L_5 . The solutions of L_5 read

$$S_0 = x - \frac{4085556703324323840000}{1644556073} x^2 + \dots \quad (70)$$

and:

$$S_1 = x^{1/5} \cdot \left(1 - \frac{80573128344696}{5} x + \frac{851459104996461085786368168}{25} x^2 + \dots \right), \\ S_2 = x^{2/5} \cdot \left(1 - \frac{3124401548255651}{465} x + \frac{9703780710544581292971588992}{775} x^2 + \dots \right), \\ S_3 = x^{3/5} \cdot \left(1 - \frac{621945576635752328}{148605} x + \frac{31428560280309440232822493239667}{4458150} x^2 \right), \\ S_4 = x^{4/5} \cdot \left(1 - \frac{2163813797006375923833}{701805760} x \right. \\ \left. + \frac{2096632093647521705592575109262587}{438628600} x^2 + \dots \right). \quad (71)$$

The series (67) can be written as a linear combination of (69) and (70) :

$$y_5 = 41113901825 \cdot (r(x) - S_0). \quad (72)$$

The series (68) can be written as a linear combination of the solutions of (71):

$$y_{1/5} = \omega \cdot S_1 - 744 \cdot \omega^2 \cdot S_2 + 356652 \cdot \omega^3 \cdot S_3 - 140361152 \cdot \omega^4 \cdot S_4 \\ + 8222780365 \cdot S_0 + 41113901825 \cdot r(x), \quad (73)$$

where $\omega^5 = 1$. The series (67) and (68) can be written respectively

$$y_5 = \tilde{X}(\tilde{Q}(x)^5) \quad \text{and:} \quad y_{1/5} = \tilde{X}(\omega \cdot \tilde{Q}(x)^{1/5}), \quad (74)$$

where $\omega^5 = 1$. They amount, respectively, to changing the nome as follows: $q \rightarrow q^5$, and its compositional inverse $q \rightarrow \omega \cdot q^{1/5}$ where $\omega^5 = 1$. These two series, (67) and (68), are actually solutions of the Schwarzian equation (40), with $W(x)$ given by (41). Note that we have the following relation:

$$\begin{aligned} 5 \cdot y_5 \cdot (1 - 1728 \cdot y_5)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y_5\right)^2 \\ = x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \cdot \frac{dy_5}{dx}, \end{aligned} \quad (75)$$

i.e.

$$5 \cdot F(y_5) = F(x) \cdot \frac{dy_5}{dx}, \quad (76)$$

and:

$$5 \cdot \frac{dx}{F(x)} = \frac{dy_5}{F(y_5)} \quad \text{and:} \quad \frac{1}{5} \cdot \frac{dx}{F(x)} = \frac{dy_{1/5}}{F(y_{1/5})}. \quad (77)$$

Remark 3.1: The series (46), (56), (68) (and also (80) below) can be seen to be functions of $\omega \cdot x^{1/N}$ with $\omega^N = 1$.

3.6. $q \rightarrow q^4$

We are not going to give explicitly the modular equation corresponding to $q \rightarrow q^4$ because it becomes a bit too large. Let us just say that it can (easily) be obtained by the elimination of t in its rational parameterization [19]:

$$x = \frac{t \cdot (t + 16)}{(t^2 + 16t + 16)^3} \quad \text{and:} \quad y = \frac{t^4 \cdot (t + 16)}{(t^2 + 256t + 4096)^3}. \quad (78)$$

This modular curve $\Gamma_4(x, y) = \Gamma_4(y, x) = 0$ can also be obtained from the elimination of the variable z between the (fundamental) modular equation $\Gamma_2(x, z) = 0$, given by (43), and the same modular equation $\Gamma_2(z, y) = 0$. The calculation of the resultant, in z , between $\Gamma_2(x, z)$ and $\Gamma_2(z, y)$ factorizes, and gives $(x - y)^2 \cdot \Gamma_4(x, y)$. This modular curve $\Gamma_4(x, y) = \Gamma_4(y, x) = 0$, has the following algebraic series solutions

$$\begin{aligned} y_4 = & x^4 + 2976x^5 + 6322896x^6 + 11838151424x^7 + 20872495228416x^8 \\ & + 35647177050980352x^9 + 59796357134115627008x^{10} \\ & + 99264875397039869263872x^{11} + \dots \end{aligned} \quad (79)$$

$$\begin{aligned} y_{1/4}(\omega, x) = & \omega \cdot x^{1/4} - 744 \cdot \omega^2 \cdot x^{1/2} + 356652 \cdot \omega^3 \cdot x^{3/4} - 140361152 \cdot x^{4/4} \\ & + 49336682376 \cdot \omega \cdot x^{5/4} - 16114625945856 \cdot \omega^2 \cdot x^{6/4} \\ & + 4999042676442272 \cdot \omega^3 \cdot x^{7/4} - 1492669488513712128 \cdot x^{8/4} \\ & + 432762805367932714848 \cdot \omega \cdot x^{9/4} + \dots \end{aligned} \quad (80)$$

where $\omega^4 = 1$, together with the (involutive) series:

$$\begin{aligned} y_1 = & -x - 1488x^2 - 2214144x^3 - 3337633792x^4 - 5094329942016x^5 \\ & - 7859077093785600x^6 - 12234039128005541888x^7 \\ & - 19190712499154486034432x^8 - 30301349938167862039412736x^9 + \dots \end{aligned} \quad (81)$$

The radius of convergence of the series (79), or (81), is $R = 1/1728$, corresponding to the vanishing of the discriminant of the modular equation $\Gamma_4(x, y) = \Gamma_4(y, x) = 0$. These three series (79), (80) and (81), can be written respectively

$$\tilde{X}(\tilde{Q}(x)^4) \quad \text{and:} \quad \tilde{X}(\omega \cdot \tilde{Q}(x)^{1/4}) \quad \text{and:} \quad \tilde{X}(-\tilde{Q}(x)), \quad (82)$$

where $\omega^4 = 1$. These series can be obtained from the series (45) and (46) of subsection (3.1). It is straightforward to see[†] that $y_4(x) = y_2(y_2(x))$, and that $y_{1/4}(x) = y_{1/2}(y_{1/2}(x))$, which amounts, *on the nome*, to performing $q \rightarrow q^2 \rightarrow (q^2)^2 = q^4$ and similarly $q \rightarrow \pm q^{1/2} \rightarrow \pm(\pm q^{1/2})^{1/2} = \omega \cdot q^{1/4}$, where $\omega^4 = 1$. However, the composition of y_2 and $y_{1/2}$ also corresponds, on the nome, to

$$q \rightarrow \pm q^{1/2} \rightarrow (\pm q^{1/2})^2 = q \quad \text{or:} \quad q \rightarrow q^2 \rightarrow \pm(q^2)^{1/2} = \pm q. \quad (83)$$

Getting rid of the identity transformation, we get $q \rightarrow -q$, which precisely corresponds to the *involution* series (81). These three series (79), (80) and (81) are actually solutions of the Schwarzian equation (40), with $W(x)$ given by (41). Note that we have the following relation:

$$\begin{aligned} & 4 \cdot y_4 \cdot (1 - 1728 \cdot y_4)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y_4\right)^2 \\ &= x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \cdot \frac{dy_4}{dx}, \end{aligned} \quad (84)$$

$$\begin{aligned} 4 \cdot \frac{dx}{F(x)} &= \frac{dy_4}{F(y_4)} \quad \text{and:} \quad \frac{1}{4} \cdot \frac{dx}{F(x)} = \frac{dy_{1/4}}{F(y_{1/4})} \\ \text{and:} \quad \frac{dx}{F(x)} &= \frac{dy_1}{F(y_1)}. \end{aligned} \quad (85)$$

3.7. Linear differential operators for $q \rightarrow q^4$

The previous algebraic series (79), (80) and (81) are solutions of an order-six linear differential operator $M_6 = M_1 \oplus M_2 \oplus M_3$, which is the *direct sum* (LCLM) of an order-three linear differential operator M_3 , an order-three linear differential operator M_2 , and an order-one linear differential operator M_1 . Let us introduce $S_i^{(n)}$ ($i = 1, \dots, n$) the (normalized) solution-series of the linear differential operators M_n ($n = 1, 2, 3$), namely the (normalized) solution of the order-one linear differential operator M_1

$$S_1^{(1)} = x + \frac{1990225984684950000}{187148203} \cdot x^2 + \frac{12420842277895932711852000000}{187148203} \cdot x^3 + \dots,$$

together with the two (normalized) solutions of the order-two linear differential operator M_2

$$\begin{aligned} S_1^{(2)} &= x + \frac{331704330780824752}{31191367} \cdot x^2 + \frac{2070140379649322118641630976}{31191367} \cdot x^3 + \dots, \\ S_2^{(2)} &= x^{1/2} \cdot \left(1 + \frac{671442747744}{31} \cdot x + \frac{5106946630014945047040}{31} \cdot x^2 + \dots\right), \end{aligned}$$

[†] The composition/iteration of *multivalued* functions, like algebraic functions, is a bit tricky, we have, however, no problem to compose *algebraic series*, for instance $x \rightarrow y_2(x) \rightarrow y_4(x) = y_2(y_2(x))$.

and the (normalized) solutions of the order-three linear differential operator M_3 :

$$\begin{aligned} S_1^{(3)} &= x + 1488x^2 + 2214144x^3 + 3337633793x^4 + 5094329944992x^5 + \dots, \\ S_2^{(3)} &= x^{1/4} \cdot \left(1 + 49336682376x + 432762805367932714848x^2 + \dots\right), \\ S_3^{(3)} &= x^{3/4} \cdot \left(1 + \frac{1249760669110568}{89163}x + \frac{2838197560249922422013408}{29721}x^2 + \dots\right). \end{aligned}$$

Note that $S_1^{(1)}$, the solution of M_1 , is a rational function:

$$S_1^{(1)} = x \cdot \frac{p_5(x)}{187148203 \cdot (1 - 2835810000x + 6549518250000x^2)^3}, \quad (86)$$

where:

$$\begin{aligned} p_5(x) &= 187148203 + 398075748036660000x + 4173268788948807866250000x^2 \\ &\quad + 62885546488332818428095703125x^3 - 31422194354407801042441406250000x^4 \\ &\quad - 121645442598919219468125000000000000x^5. \end{aligned} \quad (87)$$

Remark 3.2: The order of the linear operator M_6 , corresponds to the four series of the form $y_{1/4}$, together with the series y_4 , and the series y_1 , namely $6 = 4 + 1 + 1$. The series y_4 (given by (79)) can be seen to be an (algebraic) *analytic continuation of the involutive series* y_1 (given by (81)).

Remark 3.3: Taking into account $1 + \omega + \omega^2 + \omega^3 = 0$, let us consider the sum of the four algebraic series $y_{1/4}$. This sum reads:

$$\begin{aligned} &y_{1/4}(1, x) + y_{1/4}(\omega, x) + y_{1/4}(\omega^2, x) + y_{1/4}(\omega^3, x) \\ &= -561444608x - 5970677954054848512x^2 \\ &\quad - 37262526833687798135553785856x^3 \\ &\quad - 185766744391994261104411840078449475584x^4 \\ &\quad - 817583724079763955212555161997757454304107560960x^5 + \dots \end{aligned} \quad (88)$$

We have the following relations:

$$\begin{aligned} &y_{1/4}(1, x) + y_{1/4}(\omega, x) + y_{1/4}(\omega^2, x) + y_{1/4}(\omega^3, x) \\ &= -187148202 \cdot S_1^{(2)} - 374296406 \cdot S_1^{(1)} \end{aligned} \quad (89)$$

$$\begin{aligned} &y_{1/4}(1, x) + \omega \cdot y_{1/4}(\omega, x) + \omega^2 \cdot y_{1/4}(\omega^2, x) + \omega^3 \cdot y_{1/4}(\omega^3, x) \\ &= 1426608 \cdot S_3^{(3)}, \end{aligned} \quad (90)$$

$$\begin{aligned} &y_{1/4}(1, x) + \omega^2 \cdot y_{1/4}(\omega, x) + \omega^4 \cdot y_{1/4}(\omega^2, x) + \omega^6 \cdot y_{1/4}(\omega^3, x) \\ &= y_{1/4}(1, x) - y_{1/4}(\omega, x) + y_{1/4}(\omega^2, x) - y_{1/4}(\omega^3, x) \\ &= -2976 \cdot S_2^{(2)}, \end{aligned} \quad (91)$$

$$\begin{aligned} &y_{1/4}(1, x) + \omega^3 \cdot y_{1/4}(\omega, x) + \omega^6 \cdot y_{1/4}(\omega^2, x) + \omega^9 \cdot y_{1/4}(\omega^3, x) \\ &= 4 \cdot S_2^{(3)}. \end{aligned} \quad (92)$$

Furthermore, we have the two relations:

$$187148202 \cdot (S_1^{(2)} - S_1^{(1)}) - S_1^{(1)} = y_1 + y_4, \quad (93)$$

$$S_1^{(3)} = -y_1 + y_4. \quad (94)$$

From (89) and (93) we get

$$S_1^{(1)} = -\frac{1}{561444609} \cdot (Q_0 + y_1 + y_4), \quad (95)$$

$$S_1^{(2)} = -\frac{1}{561444606} \cdot (Q_0 - 2y_1 - 2y_4), \quad (96)$$

where:

$$Q_0 = y_{1/4}(1, x) + y_{1/4}(\omega, x) + y_{1/4}(\omega^2, x) + y_{1/4}(\omega^3, x). \quad (97)$$

We can, thus, express all the six series $S_i^{(n)}$, solutions of M_6 , in terms of the algebraic series y_1 , y_4 and $y_{1/4}(\omega^n, x)$, solutions of the modular equation $\Gamma_4(x, y) = 0$. Conversely the algebraic series y_1 , y_4 and $y_{1/4}(\omega^n, x)$ can be expressed in terms of the six series $S_i^{(n)}$. We have from (93) and (94)

$$y_1 = 93574101 \cdot (S_1^{(2)} - S_1^{(1)}) - \frac{1}{2} \cdot (S_1^{(3)} + S_1^{(1)}), \quad (98)$$

$$y_4 = 93574101 \cdot (S_1^{(2)} - S_1^{(1)}) + \frac{1}{2} \cdot (S_1^{(3)} - S_1^{(1)}). \quad (99)$$

and from (89), (90), (91), (92):

$$\begin{aligned} y_{1/4}(\omega, x) &= \omega \cdot S_2^{(3)} - 744 \cdot \omega^2 \cdot S_2^{(2)} + 356652 \cdot \omega^3 \cdot S_3^{(3)} \\ &\quad - \frac{93574101}{2} \cdot (S_1^{(2)} + 2S_1^{(1)}) - \frac{1}{2} \cdot S_1^{(1)}. \end{aligned} \quad (100)$$

The identification of the LHS of the modular equation $\Gamma_4(x, y) = 0$ with the polynomial

$$\begin{aligned} P(y) &= (y - y_1) \cdot (y - y_4) \\ &\quad \times (y - y_{1/4}(1, x)) \cdot (y - y_{1/4}(\omega, x)) \cdot (y - y_{1/4}(\omega^2, x)) \cdot (y - y_{1/4}(\omega^3, x)), \end{aligned} \quad (101)$$

gives, straightforwardly, relation (95) together with relation (86) and also:

$$\begin{aligned} &y_1 \cdot y_4 \cdot y_{1/4}(1, x) \cdot y_{1/4}(\omega, x) \cdot y_{1/4}(\omega^2, x) \cdot y_{1/4}(\omega^3, x) \\ &= \frac{1}{(1 - 2835810000x + 6549518250000x^2)^3}. \end{aligned} \quad (102)$$

Remark 3.4: The algebraic series y_1 , y_4 , $y_{1/4}(\omega^n, x)$, solutions of the modular equation $\Gamma_4(x, y) = 0$, can be expressed as linear combinations of the solutions of the three linear differential operators M_n , $n = 1, 2, 3$. If one introduces the (finite) Galois group of the polynomial associated with the modular equation $\Gamma_4(x, y) = 0$, and the differential Galois groups of the three linear differential operators M_n , one sees that the relation between these different Galois groups is far from being straightforward.

3.8. More correspondence series

Let us display ¶ more correspondence series. More examples of correspondence series are displayed in Appendix C.

¶ For all these examples we used gfun of Bruno Salvy. We used the following commands: `algeqtoDIFFeq`, `diffEQtohomDIFFeq`, `de2DIFFop`, `algeqtoseries`, `formal_sols`.

- The (algebraic) series

$$\begin{aligned} \tilde{X}(\tilde{Q}(x)^5) = & x^5 + 3720x^6 + 9287460x^7 + 19648405600x^8 + 38124922672650x^9 \\ & + 70330386411705000x^{10} + \dots \end{aligned} \quad (103)$$

is solution of a modular equation $\Gamma_5(x, y) = \Gamma_5(y, x) = 0$, that we will not write here, but can easily be obtained from its rational parametrization [19]:

$$x = \frac{t}{(t^2 + 10t + 5)^3}, \quad y = \frac{t^5}{(t^2 + 250t + 3125)^3}. \quad (104)$$

This series (103) is solution of an order-six linear differential operator $L_6 = L_1 \oplus L_5$, which is the direct sum of an order-one linear differential operator L_1 with a rational function solution (69), and an irreducible order-five linear differential operator L_5 .

- The (algebraic) series

$$\begin{aligned} \tilde{X}(\tilde{Q}(x)^6) = & x^6 + 4464x^7 + 12805560x^8 + 30222607872x^9 \\ & + 64062187946172x^{10} + \dots \end{aligned} \quad (105)$$

is solution of a modular equation $\Gamma_6(x, y) = \Gamma_6(y, x) = 0$, that will not be written here, but can easily be obtained from its rational parametrization [19]:

$$\begin{aligned} x = & \frac{t \cdot (t+8)^3 \cdot (t+9)^2}{(t+6)^3 \cdot (t^3 + 18t^2 + 84t + 24)^3}, \\ y = & \frac{t^6 \cdot (t+8)^2 \cdot (t+9)^3}{(t+12)^3 \cdot (t^3 + 252t^2 + 3888t + 15552)^3}. \end{aligned} \quad (106)$$

This series (105) is solution of an order-twelve linear differential operator $L_{12} = L_1 \oplus L_{11}$, which is the *direct sum* of an order-one linear differential operator L_1 with a rational function solution of the form

$$x \cdot \frac{p_{11}(x)}{(54000x - 1)^3 \cdot q_3(x)^3}, \quad (107)$$

where $p_{11}(x)$ is a polynomial of degree eleven, and where $q_3(x)$ reads

$$187999470568800000000x^3 - 224179462188000000x^2 + 151013228706000x - 1,$$

and an order-eleven linear differential operator L_{11} .

- We can also consider

$$\tilde{X}(\tilde{Q}(x)^{13}) = x^{13} + 9672x^{14} + 52931268x^{15} + 216226356320x^{16} + \dots \quad (108)$$

which is solution of a modular equation[†] $\Gamma_{13}(x, y) = \Gamma_{13}(y, x) = 0$, that we will not write here, but can easily be obtained from its rational parametrization [19]:

$$\begin{aligned} x = & \frac{t}{(t^2 + 5t + 13) \cdot (t^4 + 7t^3 + 20t^2 + 19t + 1)^3}, \\ y = & \frac{t^{13}}{(t^2 + 5t + 13) \cdot (t^4 + 247t^3 + 3380t^2 + 15379t + 28561)^3}. \end{aligned} \quad (109)$$

This series (108) is solution of an order-fourteen linear differential operator $L_{14} = L_1 \oplus L_{13}$, which is the direct sum of an order-one linear differential operator L_1 with a rational function solution, and an irreducible order-thirteen linear differential operator L_{13} .

[†] The polynomial $\Gamma_{13}(x, y)$ is of degree 14 in y (or x).

- Let us consider

$$\tilde{X}(\tilde{Q}(x)^9) = x^9 + 6696x^{10} + 26681076x^{11} + 82647211104x^{12} + \dots \quad (110)$$

which is solution of a modular equation $\Gamma_9(x, y) = \Gamma_9(y, x) = 0$, that we will not write here, but can easily be obtained from its rational parametrization [19]:

$$x = \frac{t \cdot (t^2 + 9t + 27)}{(t+3)^3 \cdot (t^3 + 9t^2 + 27t + 3)^3}, \quad y = \frac{t^9 \cdot (t^2 + 9t + 27)}{(t+9)^3 \cdot (t^3 + 243t^2 + 2187t + 6561)^3}.$$

The polynomial $\Gamma_9(x, y)$ is of degree 12 in y (resp. in x). We thus have twelve algebraic solutions-series of the *modular equation* $\Gamma_9(x, y) = 0$. This series (110) is solution of an order-twelve linear differential operator $L_{12} = L_1 \oplus L_{11}$, which is the direct sum of an order-one operator L_1 with a rational function solution of the form

$$x \cdot \frac{q_{11}(x)}{(12288000x + 1)^3 \cdot q_3(x)^3}, \quad (111)$$

where $q_{11}(x)$ is a polynomial of degree eleven, and where $q_3(x)$ reads

$$\begin{aligned} & 3338586724673519616000000000x^3 - 3750657365033091072000000x^2 \\ & + 1855762905734664192000x + 1, \end{aligned} \quad (112)$$

and an order-eleven linear differential operator L_{11} . The (nine) series which are compositional inverse of the series (110), are also solutions of the modular equation $\Gamma_9(x, y) = 0$, read:

$$\begin{aligned} \tilde{X}(\tilde{Q}(x)^{1/9}) &= \omega \cdot x^{1/9} - 744 \cdot \omega^2 \cdot x^{2/9} + 356652 \cdot \omega^3 \cdot x^{1/3} \\ &- 140361152 \cdot \omega^4 \cdot x^{4/9} + 49336682190 \cdot \omega^5 \cdot x^{5/9} - 16114625669088 \cdot \omega^6 \cdot x^{2/3} \\ &+ 4999042477430456 \cdot \omega^7 \cdot x^{7/9} + \dots \end{aligned} \quad (113)$$

where $\omega^9 = 1$. These (nine) series (113) are solutions of the order-twelve linear differential operator L_{12} . Note that the (two) *order-three* series

$$\begin{aligned} y_\omega(x) &= y_{1/3}(y_3(x)) = \omega \cdot x - 744 \cdot \omega \cdot (\omega - 1) \cdot x^2 \\ &+ 36 \cdot \omega \cdot (\omega - 1) \cdot (9907\omega - 20845) \cdot x^3 \\ &- 32 \cdot \omega \cdot (\omega - 1) \cdot (-24876477\omega + 22887765) \cdot x^4 + \dots \end{aligned} \quad (114)$$

where $\omega^2 + \omega + 1 = 0$, are also solutions of the modular equation $\Gamma_9(x, y) = 0$, and also of the order-twelve operator L_{12} . We thus have $1 + 2 + 9 = 12$ *algebraic* solutions of the modular equation $\Gamma_9(x, y) = 0$, and solutions of L_{12} .

- The (algebraic) series

$$\tilde{X}(\tilde{Q}(x)^{10}) = x^{10} + 7440x^{11} + 32413320x^{12} + 108395513600x^{13} + \dots \quad (115)$$

is solution of a modular equation $\Gamma_{10}(x, y) = \Gamma_{10}(y, x) = 0$, which has the rational parameterization [19]:

$$\begin{aligned} x &= \frac{t \cdot (t+4)^5 \cdot (t+5)^2}{(t^6 + 20t^5 + 160t^4 + 640t^3 + 1280t^2 + 1040t + 80)^3} \\ y &= \frac{t^{10} \cdot (t+4)^2 \cdot (t+5)^5}{(t^6 + 260t^5 + 6400t^4 + 64000t^3 + 320000t^2 + 800000t + 800000)^3}. \end{aligned} \quad (116)$$

The degree of the polynomial in $\Gamma_{10}(x, y) = 0$ in y (resp. in x) is 18. The other (algebraic) series solutions of $\Gamma_{10}(x, y) = 0$ are the compositional inverse of series (115), namely

$$\begin{aligned} y_{1/10}(x) = & \omega \cdot x^{1/10} - 744 \cdot \omega^2 \cdot x^{2/10} + 356652 \cdot \omega^3 \cdot x^{3/10} \\ & - 140361152 \cdot \omega^4 \cdot x^{4/10} + 49336682190 \cdot \omega^5 \cdot x^{5/10} \\ & - 16114625669088 \cdot \omega^6 \cdot x^{6/10} + \dots \end{aligned} \quad (117)$$

where $\omega^{10} = 1$, together with[†]

$$\begin{aligned} y_{5/2}(x) = & \omega \cdot x^{5/2} + 1860 \cdot \omega \cdot x^{7/2} + 2913930 \cdot \omega \cdot x^{9/2} - 744 \cdot x^{10/2} \\ & + 4404293000 \cdot \omega \cdot x^{11/2} - 2767680 \cdot x^6 + 6624982333875 \cdot \omega \cdot x^{13/2} + \dots \end{aligned} \quad (118)$$

where $\omega^2 = 1$, and

$$\begin{aligned} y_{2/5}(x) = & \omega \cdot x^{2/5} - 744 \cdot \omega^2 \cdot x^{4/5} + 356652 \cdot \omega^3 \cdot x^{6/5} + \frac{1488}{5} \cdot \omega \cdot x^{7/5} \\ & - 140361152 \cdot \omega^4 \cdot x^{8/5} - \frac{2214144}{5} \cdot \omega^2 \cdot x^{9/5} + 49336682190 \cdot \omega^5 \cdot x^{10/5} \\ & + \frac{1592094528}{5} \cdot \omega^3 \cdot x^{11/5} + \dots \end{aligned} \quad (119)$$

where $\omega^5 = 1$. We thus have $1 + 2 + 5 + 10 = 18$ algebraic solutions of $\Gamma_{10}(x, y) = 0$ and of L_{18} . The order-eighteen linear differential operator L_{18} is the *direct sum* of an order-seventeen linear differential operator L_{17} , and an order-one linear differential operator L_1 , which has a rational function solution,

$$x \cdot \frac{p_{17}}{p_6^3}, \quad (120)$$

where p_{17} is a polynomial of degree 17, where p_6 reads

$$\begin{aligned} & 66661978554978958501295319312489107870472732672000 x^6 \\ & + 62082816308629282586712746552975312469884928000 x^5 \\ & + 21122955530832902270001123584504233628467200 x^4 \\ & - 233405320133674124312518469774131200 x^3 \\ & + 32278855882815402576742692253440 x^2 \\ & - 428244362959801779810720 x + 1. \end{aligned} \quad (121)$$

• The (algebraic) series

$$\tilde{X}(\tilde{Q}(x)^{25}) = x^{25} + 18600 x^{26} + 184821300 x^{27} + 1304017532000 x^{28} + \dots \quad (122)$$

is solution of a *modular equation* $\Gamma_{25}(x, y) = \Gamma_{25}(y, x) = 0$, that we will not write here, but can easily be obtained from its rational parametrization [19]

$$x = t \cdot \frac{p_5}{p_{10}^3}, \quad y = t^{25} \cdot \frac{p_5}{q_{10}^3}, \quad (123)$$

where:

$$p_5 = t^4 + 5t^3 + 15t^2 + 25t + 25, \quad (124)$$

$$\begin{aligned} p_{10} = & t^{10} + 10t^9 + 55t^8 + 200t^7 + 525t^6 + 1010t^5 + 1425t^4 \\ & + 1400t^3 + 875t^2 + 250t + 5, \end{aligned} \quad (125)$$

$$\begin{aligned} q_{10} = & t^{10} + 250t^9 + 4375t^8 + 35000t^7 + 178125t^6 + 631250t^5 \\ & + 1640625t^4 + 3125000t^3 + 4296875t^2 + 3906250t + 1953125. \end{aligned} \quad (126)$$

[†] The series (118) corresponds to $y_{1/2}(y_5(x))$.

The polynomial in the modular equation $\Gamma_{25}(x, y) = 0$ is of degree 30 in y (resp. in x), and thus has thirty algebraic solution series, corresponding to the series (122), together with the 25 compositional inverse of series (122), namely

$$\begin{aligned} \omega \cdot x^{1/25} &- 744 \cdot \omega^2 \cdot x^{2/25} + 356652 \cdot \omega^3 \cdot x^{3/25} - 140361152 \cdot \omega^4 \cdot x^{4/25} \\ &+ 49336682190 \cdot \omega^5 \cdot x^{5/25} + \dots \end{aligned} \quad (127)$$

where $\omega^{25} = 1$, together with the four (order-five[†]) series

$$y_{1/5}(y_5(x)) = \omega \cdot x - 744 \cdot \omega \cdot (\omega - 1) \cdot x^2 + \dots \quad (128)$$

with $\omega^5 = 1$ but $\omega \neq 1$. This thus gives $30 = 1 + 4 + 25$ algebraic series. They are solutions of an order-30 linear differential operator which is the direct-sum of an order-29 linear differential operator, and an order-one linear differential operator with a rational function solution

$$x \cdot \frac{p_{29}}{p_{10}^3}, \quad (129)$$

where p_{29} is a polynomial of degree 29, and where p_{10} is a polynomial of degree 10.

4. The one-parameter series solutions of the Schwarzian equation.

The Schwarzian equation (40) has more solutions than the infinite discrete set of algebraic series (see (45), (55), (67), (79), (105), (108), ...) corresponding to *modular correspondences*. One also has a series *depending on one parameter*, namely:

$$\begin{aligned} y(a, x) = & a \cdot x - 744 \cdot a \cdot (a - 1) \cdot x^2 + 36 \cdot a \cdot (a - 1) \cdot (9907a - 20845) \cdot x^3 \\ & - 32 \cdot a \cdot (a - 1) \cdot (4386286a^2 - 20490191a + 27274051) \cdot x^4 \\ & + 6 \cdot a \cdot (a - 1) \cdot (8222780365a^3 - 61396351027a^2 \\ & + 171132906629a - 183775457147) \cdot x^5 \\ & - 144 \cdot a \cdot (a - 1) \cdot (111907122702a^4 - 1162623833873a^3 + 5000493989295a^2 \\ & - 10801207072185a + 10212230113145) \cdot x^6 \\ & + 8 \cdot a \cdot (a - 1) \cdot (624880309678807a^5 - 8367080813672297a^4 \\ & + 48909476982869878a^3 - 158792594445015178a^2 \\ & + 293243568886999823a - 254689844062110385) \cdot x^7 \\ & - 192 \cdot a \cdot (a - 1) \cdot (7774319708776120a^6 - 127824707491524999a^5 \\ & + 946950323149342341a^4 - 4101941044701784034a^3 \\ & + 11156847890086765926a^2 - 18508096006772656203a \\ & + 15126379507970624425) \cdot x^8 + \dots \end{aligned} \quad (130)$$

Note that all the algebraic series (81), (114), (128), (see also (152) below), ... associated with modular equations, are of the form (130) where the parameter is a N -th root of unity: $a^N = 1$.

Note that this one-parameter series (130) is a series of the form

$$y(a, x) = a \cdot x + a \cdot (a - 1) \cdot \sum_{n=2}^{\infty} P_n(a) \cdot x^n, \quad (131)$$

[†] The composition of series (128) with itself five times gives the identity transformation.

where the polynomials $P_n(a)$ are polynomials of degree $n - 2$ in the parameter a , with integer coefficients[‡].

This one-parameter series (130), (131) verifies the following composition rule:

$$y(a, y(a', x)) = y(a', y(a, x)) = y(a a', x). \quad (132)$$

These series commute. One can verify that this one-parameter series (130) can, in fact, be written

$$y(a, x) = \tilde{X}(a \cdot \tilde{Q}(x)), \quad (133)$$

where

$$\begin{aligned} \tilde{X}(q) = & q - 744 q^2 + 356652 q^3 - 140361152 q^4 + 49336682190 q^5 \\ & - 16114625669088 q^6 + 4999042477430456 q^7 + \dots \end{aligned} \quad (134)$$

and[†] its composition inverse:

$$\begin{aligned} \tilde{Q}(x) = & x + 744 x^2 + 750420 x^3 + 872769632 x^4 + 1102652742882 x^5 \\ & + 1470561136292880 x^6 + 2037518752496883080 x^7 + \dots \end{aligned} \quad (135)$$

The nome series (135) has a radius of convergence $R = 1/1728 = 0.00057870370 \dots$

In the $a \rightarrow 0$ limit one has

$$\begin{aligned} \lim_{a \rightarrow 0} \frac{y(a, x)}{a} = & x + 744 x^2 + 750420 x^3 + 872769632 x^4 + 1102652742882 x^5 \\ & + 1470561136292880 x^6 + 2037518752496883080 x^7 + 2904264865530359889600 x^8 \\ & + 4231393254051181981976079 x^9 + \dots \end{aligned} \quad (136)$$

which is nothing but the nome series $\tilde{Q}(x)$ given by (135). In the $a \rightarrow \infty$ limit one has

$$\begin{aligned} \lim_{a \rightarrow \infty} y\left(a, \frac{x}{a}\right) = & x - 744 x^2 + 356652 x^3 - 140361152 x^4 + 49336682190 x^5 \\ & - 16114625669088 x^6 + 4999042477430456 x^7 - 1492669384085015040 x^8 \\ & + 432762759484818142437 x^9 + \dots \end{aligned} \quad (137)$$

which is nothing but \tilde{X} , the (elliptic modulus) series (134).

Let us introduce the ratio of the polynomials in expansion (131):

$$R_n(a) = \frac{P_n(a)}{P_{n+1}(a)}. \quad (138)$$

One finds, in the $n \rightarrow \infty$ and $a \rightarrow 0$ limit, that the ratio (138) becomes $1/1728 = 0.00057870 \dots$. For miscellaneous small values of the parameter a one can see, that this ratio (138) also becomes $1/1728$ in the $n \rightarrow \infty$ limit.

In the last $n \rightarrow \infty$ and $a \rightarrow \infty$ limit (137), the ratio (138) becomes^{††} $-0.004316810242 \dots$ which corresponds to the radius of convergence of the series

[‡] This can be seen as a consequence of the fact that $y(a, x) = \tilde{X}(a \cdot \tilde{Q}(x))$, where $\tilde{X}(x)$ and $\tilde{Q}(x)$ are actually series with integer coefficients (see (22) and (23)).

[†] In Maple the $\tilde{X}(q)$ series (22), (134) can be obtained substituting $L = \text{EllipticModulus}(q^{1/2})^2$, in $1/j = L^2 \cdot (L-1)^2/(L^2 - L + 1)^3/256$. See <https://oeis.org/A066395> for the series (22) and <https://oeis.org/A091406> for the series (23).

^{††} Obtained with 421 coefficients.

(22), (134). This radius of convergence is according to Vaclav Kotesovec¶

$$\exp\left(-\sqrt{3} \cdot \pi\right) = 0.004333420501 \dots \quad (139)$$

which is reminiscent of the selected values (see equation (55) in [32]):

$$t = \exp\left(i\pi \frac{1+i\sqrt{3}}{2}\right) = i \cdot \exp\left(-\frac{\sqrt{3}}{2} \cdot \pi\right) \quad \text{or:} \quad j\left(\frac{1+i\sqrt{3}}{2}\right) = 0. \quad (140)$$

The nearest to $x = 0$ singularity of \tilde{X} is thus $x_c = t^2 = -\exp(-\sqrt{3} \cdot \pi)$. We have seen that the radius of convergence of the series (81) (i.e. $a = -1$) is $R = 1/1728$, corresponding to the vanishing of the discriminant of the modular equation $\Gamma_4(x, y) = \Gamma_4(y, x) = 0$, and more generally, for $|a| = 1$, one can see that the radius of convergence of the series (130), (131) for N -th root of unity, $a^N = 1$, is also‡ $R = 1/1728$.

More generally, the radius of convergence of (130), (131) corresponds to the singularities of (133), namely the $x = 1/1728$ singularity of $\tilde{Q}(x)$, and to the values of x such that $a \cdot \tilde{Q}(x) = -\exp(-\sqrt{3} \cdot \pi)$, which corresponds to the singularity of $\tilde{X}(x)$, namely:

$$x = \tilde{X}\left(-\frac{1}{a} \cdot \exp(-\sqrt{3} \cdot \pi)\right). \quad (141)$$

When the parameter a is large enough ($|a| > \simeq 7.5$), the radius of convergence no longer corresponds to $R = 1/1728$, but to the singularity (141).

This transcendental value (139), for the radius of convergence of the series $\tilde{X}(q)$, is a strong incentive to understand the “very nature” of the one-parameter series (130), (131), especially since it can be written in the simple form (133). Generically the one-parameter series (130), being solution of a Schwarzian equation, is a *differentially algebraic series*, but is it possible that this series could be, only for *selected values* of the parameter, an algebraic series, or just a D -finite series, or possibly a D - D -finite series ?

5. Trying to understand the one-parameter series solutions.

5.1. When the one-parameter series becomes an algebraic series

For $a = -1$ the (*involution*) series $y(a, x)$ (see series (81))

$$\begin{aligned} -x &- 1488x^2 - 2214144x^3 - 3337633792x^4 - 5094329942016x^5 \\ &- 7859077093785600x^6 - 12234039128005541888x^7 + \dots \end{aligned} \quad (142)$$

has a radius of convergence $1/1728 = 0.00057870\dots$. Let us generalize what we have seen in subsection (3.6) with series (81). Let us first recall the algebraic series (corresponding to $q \rightarrow q^3$) y_3 , given by (55), and $y_{1/3}$, given by (56), where $\omega^3 = 1$, and combine y_3 and $y_{1/3}$. We first get:

$$y_3\left(y_{1/3}(x)\right) = x. \quad (143)$$

¶ See <https://oeis.org/A066395> and <https://oeis.org/A066395/b066395.txt> for the reciprocal of j -function. See also in [81], $Q(\exp(-\sqrt{3} \cdot \pi)) = 0$ or $J(\exp(-\sqrt{3} \cdot \pi)) = 0$, where Q is the Eisenstein series E_4 and J is the Klein modular invariant.

‡ This also corresponds to vanishing of the discriminant of the corresponding modular equations.

More interestingly we also get the following algebraic series (see (114) previously):

$$\begin{aligned} y_\omega(x) = y_{1/3}(y_3(x)) = & \omega \cdot x - 744 \cdot \omega \cdot (\omega - 1) \cdot x^2 \\ & + 36 \cdot \omega \cdot (\omega - 1) \cdot (9907\omega - 20845) \cdot x^3 \\ & - 32 \cdot \omega \cdot (\omega - 1) \cdot (22887765 - 24876477\omega) \cdot x^4 + \dots \end{aligned} \quad (144)$$

where $\omega^3 = 1$. One can verify that series (144) is actually series (130) when $a^3 = 1$. One can verify that this series is (for $\omega \neq 1$) a series of order 3:

$$y_\omega(y_\omega(y_\omega(x))) = x. \quad (145)$$

Let us also recall the algebraic series (corresponding to $q \rightarrow q^5$) y_5 , given by (67), and its compositional inverse $y_{1/5}$, given by (68), where $\omega^5 = 1$, and let us compose y_5 and $y_{1/5}$. We first get:

$$y_5(y_{1/5}(x)) = x. \quad (146)$$

More interestingly, we also get the following series (see series (128) previously):

$$\begin{aligned} y_\omega(x) = y_{1/5}(y_5(x)) = & \omega \cdot x - 744 \cdot \omega \cdot (\omega - 1) \cdot x^2 \\ & + 36 \cdot \omega \cdot (\omega - 1) \cdot (9907\omega - 20845) \cdot x^3 \\ & - 32 \cdot \omega \cdot (\omega - 1) \cdot (4386286\omega^2 - 20490191\omega + 27274051) \cdot x^4 \\ & + 6 \cdot \omega \cdot (\omega - 1) \cdot (8222780365\omega^3 - 61396351027\omega^2 \\ & \quad + 171132906629\omega - 183775457147) \cdot x^5 \\ & - 144 \cdot \omega \cdot (\omega - 1) \cdot (-1274530956575\omega^3 + 4888586866593\omega^2 \\ & \quad - 10913114194887\omega + 10100322990443) \cdot x^6 + \dots \end{aligned} \quad (147)$$

where $\omega^5 = 1$.

One can verify that (147) is actually (130) when $a^5 = 1$. One can verify that this series is (for $\omega \neq 1$) a series of order 5:

$$y_\omega(y_\omega(y_\omega(y_\omega(y_\omega(x))))) = x. \quad (148)$$

This is a straight consequence of (133) with $a^5 = 1$. Similarly, let us now consider

$$\begin{aligned} y_{13} = \tilde{X}(\tilde{Q}(x)^{13}) = & x^{13} + 9672x^{14} + 52931268x^{15} + 216226356320x^{16} \\ & + 735033166074714x^{17} + 2200510278533887632x^{18} + \dots \end{aligned} \quad (149)$$

Its compositional inverse (Puisseux) series reads

$$\begin{aligned} y_{1/13} = \tilde{X}(\tilde{Q}(x)^{1/13}) = & \omega \cdot x^{1/13} - 744\omega^2 \cdot x^{2/13} + 356652 \cdot \omega^3 \cdot x^{3/13} \\ & - 140361152 \cdot \omega^4 \cdot x^{4/13} + 49336682190 \cdot \omega^5 \cdot x^{5/13} \\ & - 16114625669088 \cdot \omega^6 \cdot x^{6/13} + \dots \end{aligned} \quad (150)$$

where $\omega^{13} = 1$. Let us compose y_{13} and $y_{1/13}$. We first get

$$y_{13}(y_{1/13}(x)) = x, \quad (151)$$

which corresponds to: $q \longrightarrow \omega q^{1/13} \longrightarrow (\omega q^{1/13})^{13} = q$.

One can also compose such an algebraic (modular correspondence) series (159) with the M -th root algebraic series (156) (here $a^M = 1$), to get more (modular correspondence) *algebraic* series:

$$y = a \cdot x^N + 744 \cdot N \cdot a \cdot x^{N+1} + \dots \quad \text{with:} \quad a^M = 1. \quad (160)$$

Series (160) is a (modular correspondence) series solution of the modular equation $\Gamma_{N \cdot M^2}(x, y) = 0$. The series (160) corresponds, in the nome, to transformation $q \rightarrow q^N \rightarrow \omega \cdot ((q^N)^M)^{1/M} = \omega \cdot q^N$, where $\omega^M = 1$.

5.1.2. The one-parameter series (130) is not generically a D -finite series

The one-parameter series (130) becomes an *algebraic series* when the parameter is a N -th root of unity. All the previous *algebraic series* associated with *modular equations*, can also be seen as D -finite series as displayed in the previous section (3.7). Along this line it is crucial to note that these series are solutions of a linear differential operator (like M_3 in the previous section (3.7)) of order increasing with N . Therefore, we see that *one cannot expect the one-parameter series (130) to be generically D -finite*, being solution a finite order linear differential operator with coefficients polynomial in x and in the parameter a , since the order of this linear differential operator *grows with N* when the parameter is a N -th root of unity.

5.2. When the one-parameter series becomes a globally bounded series

Note that, for *integer* values of the parameter a , the series $y(a, x)$ are series with *integer* coefficients. More generally, one can see easily that *such series are globally bounded* [43, 44] for any rational number $a = P/Q$: the series (130) can be recast into a series with *integer* coefficients if one rescales x as follows: $x \rightarrow Q \cdot x$.

If one of these series is D -finite, the series should be, according to Christol's conjecture [82], a diagonal of a rational (or algebraic) function [43]. In particular this series should *reduce to algebraic function modulo any prime number* [43, 44]. Let us focus, for instance, on the particular value $a = 3$. For $a = 3$ the series $y(a, x)$ is a series with *integer* coefficients

$$S = 3x - 4464x^2 + 1917216x^3 - 1013769984x^4 - 33437759328x^5 - 420498625999104x^6 - 452363497164804864x^7 + \dots \quad (161)$$

which has a radius of convergence $1/1728 = 0.00057870 \dots$. If one considers the series (161) modulo different primes p , it is very difficult to see (for p large enough) if this series (161) is an *algebraic series* modulo p , or, even, is D -finite modulo p . We have, however, found the following result. Introducing

$$\sigma = \frac{S - 3x}{3 \cdot 2^5 \cdot x} + \frac{99}{2} \cdot x + 1 = 1 + 3x + 19971x^2 - 10560104x^3 - 348309993x^4 - 4380194020824x^5 - 4712119762133384x^6 + \dots \quad (162)$$

this series reduces, modulo $p = 2$, to the algebraic series

$$\sigma(x) = 1 + x + x^2 + x^4 + x^8 + x^{16} + x^{32} + x^{64} + x^{128} + x^{256} + \dots \quad (163)$$

solution, modulo $p = 2$, of the algebraic polynomial:

$$\sigma(x^2) - \sigma(x) + x = \sigma(x)^2 - \sigma(x) + x = 0. \quad (164)$$

The nature of the series (161), or more generally of (130) for integer, or rational values of the parameter a , remains an open question. It seems that such globally bounded series are not D -finite. At least, one has an *infinite number of differentially algebraic series*. Are these globally bounded series D - D -finite series [48, 49] ?

5.3. Miscellaneous calculations.

The nome series (135) and the mirror map series (134), are, respectively, solutions of the following Schwarzian equations

$$\{\tilde{Q}(x), x\} + \frac{1}{2 \cdot \tilde{Q}(x)^2} \cdot \left(\frac{d\tilde{Q}(x)}{dx} \right)^2 + W(x) = 0, \quad (165)$$

and

$$\{\tilde{X}(x), x\} - \frac{1}{2 \cdot x^2} - W(\tilde{X}(x)) \cdot \left(\frac{d\tilde{X}(x)}{dx} \right)^2 = 0, \quad (166)$$

where:

$$W(x) = -\frac{1}{2} \cdot \frac{1 - 1968x + 2654208x^2}{x^2 \cdot (1 - 1728x)^2}. \quad (167)$$

Let us introduce the hypergeometric function:

$$F(x) = x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2. \quad (168)$$

Note that the Schwarzian equation (165), on $\tilde{Q}(x)$, can be seen to be a consequence of (see (301) below):

$$F(x) = \frac{\tilde{Q}(x)}{\tilde{Q}(x)'} \quad \text{together with:} \quad W(x) = \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)} \right)^2. \quad (169)$$

Therefore the nome $\tilde{Q}(x)$ is *also* solution of the order-one linear differential operator:

$$\mathcal{L}_1 = F(x) \cdot D_x - 1 \quad \text{where:} \quad (170)$$

$$F(x) = x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2.$$

It is thus DD -finite[†]:

$$\frac{\tilde{Q}(x)'}{\tilde{Q}(x)} = \frac{1}{F(x)} \quad \text{or:} \quad \tilde{Q}(x) = \exp\left(\int^x \frac{dx}{F(x)}\right). \quad (171)$$

The one-parameter series $y(x) = y(a, x)$, given by (130), is solution of the rank-two equation (see (32))

$$A_R(x) - A_R(y(x)) \cdot y'(x) + \frac{y''(x)}{y'(x)} = 0, \quad (172)$$

with

$$A_R(x) = \frac{F'(x)}{F(x)}, \quad (173)$$

and also solution of the Schwarzian condition

$$W(x) - W(y(x)) \cdot y'(x)^2 + \{y(x), x\} = 0, \quad (174)$$

where:

$$\begin{aligned} W(x) &= \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)} \right)^2 = A'_R(x) + \frac{A_R(x)^2}{2} \\ &= -\frac{1}{2} \cdot \frac{1 - 1968x + 2654208x^2}{x^2 \cdot (1 - 1728x)^2}. \end{aligned} \quad (175)$$

[†] See [48, 49].

Note that $W(x)$ is a *rational function*, but this is far from being the case for $A_R(x)$. We will see, in the following, that the one-parameter series $y(x) = y(a, x)$, given by (130), is also solution of:

$$a \cdot \frac{\partial y(a, x)}{\partial a} = F(y(a, x)) = F(x) \cdot \frac{\partial y(a, x)}{\partial x}. \quad (176)$$

5.4. More one-parameter series solutions.

If one combines y_2 , the “correspondence” series (45) solution of the modular equation (43), with the one-parameter series (130), one gets a one-parameter series

$$\begin{aligned} y_2^{(a)} &= y(a, y_2) = \tilde{X}\left(a \cdot \tilde{Q}(x)^2\right) = a \cdot x^2 + 1488 \cdot a \cdot x^3 \\ &\quad - 24 \cdot a \cdot (31a - 85599) \cdot x^4 - 256 \cdot a \cdot (8649a - 11180329) \cdot x^5 \\ &\quad + 12 \cdot a \cdot (29721a^2 - 392019552a + 338926406215) \cdot x^6 \\ &\quad + 192 \cdot a \cdot (8292159a^2 - 45872836768a + 30686235044193) \cdot x^7 + \dots \end{aligned} \quad (177)$$

This series (177) is also solution of the Schwarzian equation (42). Furthermore we have:

$$\begin{aligned} 2 \cdot y_2^{(a)} \cdot (1 - 1728 \cdot y_2^{(a)})^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y_2^{(a)}\right)^2 \\ = x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \cdot \frac{dy_2^{(a)}}{dx}. \end{aligned} \quad (178)$$

When $a = 1$, the radius of convergence of (177) is $1/1728 = 0.000578703703 \dots$, and this is also the case for any a , N -th root of unity $a^N = 1$. Similarly to what has been sketched in section (4) (see equation (141)), let us remark that the *one-parameter* series (177) can be written $\tilde{X}\left(a \cdot \tilde{Q}(x)^2\right)$. For generic value of the parameter a , the radius of convergence of (177) will correspond, for a small enough, to the singularity of $\tilde{Q}(x)$, namely $1/1728$, and for a large enough, to the values of x such that $a \cdot \tilde{Q}(x)^2 = -\exp(-\sqrt{3} \cdot \pi)$, which correspond to the singularity of $\tilde{X}(x)$, namely:

$$x = \tilde{X}\left(\left(-\frac{1}{a} \cdot \exp(-\sqrt{3} \cdot \pi)\right)^{1/2}\right). \quad (179)$$

More generally, all the series

$$\tilde{X}\left(a \cdot \tilde{Q}(x)^n\right) = a \cdot x^n + \dots \quad (180)$$

have a radius of convergence corresponding, for a small enough, to the occurrence of the singularity of the nome-like series $\tilde{Q}(x)$, namely $x = 1/1728$.

Similarly to (177), if one combines y_3 , the “correspondence” series (55) solution of the modular equation (53), with the one-parameter series (130), one gets a one-parameter series

$$\begin{aligned} y_3^{(a)} &= y(a, y_3) = \tilde{X}\left(a \cdot \tilde{Q}(x)^3\right) = a \cdot x^3 + 2232 \cdot a \cdot x^4 + 3911868 \cdot a \cdot x^5 \\ &\quad - 24 \cdot a \cdot (31a - 265833940) \cdot x^6 - 54 \cdot a \cdot (61504a - 187769367601) \cdot x^7 \\ &\quad - 1296 \cdot a \cdot (7351340a - 12322394107529) \cdot x^8 + \dots \end{aligned} \quad (181)$$

This series (181) is also solution of the Schwarzian equation (42). Furthermore we have:

$$\begin{aligned} & 3 \cdot y_3^{(a)} \cdot (1 - 1728 \cdot y_3^{(a)})^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y_3^{(a)}\right)^2 \\ &= x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \cdot \frac{dy_3^{(a)}}{dx}. \end{aligned} \quad (182)$$

Similarly:

$$\begin{aligned} y_5^{(a)}(x) &= y(a, y_5(x)) = \tilde{X}(a \cdot \tilde{Q}(x)^5) = a \cdot x^5 + 3720 \cdot a \cdot x^6 \\ &+ 9287460 \cdot a \cdot x^7 + 19648405600 \cdot a \cdot x^8 + 38124922672650 \cdot a \cdot x^9 \\ &- 24 \cdot a \cdot (31a - 2930432767154406) \cdot x^{10} \\ &- 40 \cdot a \cdot (138384a - 3142471028063763509) \cdot x^{11} \\ &- 960 \cdot a \cdot (25120323a - 229208433006295134073) \cdot x^{12} + \dots \end{aligned} \quad (183)$$

This series (183) is also solution of the Schwarzian equation (42). Furthermore we have:

$$\begin{aligned} & 5 \cdot y_5^{(a)} \cdot (1 - 1728 \cdot y_5^{(a)})^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot y_5^{(a)}\right)^2 \\ &= x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2 \cdot \frac{dy_5^{(a)}}{dx}. \end{aligned} \quad (184)$$

One also easily gets:

$$5 \cdot F(y_5^{(a)}(x)) = F(x) \cdot \frac{dy_5^{(a)}(x)}{dx} = 5 \cdot a \cdot \frac{\partial y_5^{(a)}(x)}{\partial a}. \quad (185)$$

More generally, let us introduce the modular correspondence series $y_n(x) = x^n + 744 \cdot n \cdot x^{n+1} + \dots$ (for $n \geq 2$), one can verify *that these series commute*. These modular correspondences $y_n(x)$ can easily be generalized to *one-parameter* series $y(a, y_n(x))$ which are also solutions of the Schwarzian equations:

$$y(a, y_n(x)) = a \cdot x^n + 744 \cdot n \cdot a \cdot x^{n+1} + \dots \quad (186)$$

Let us recall the one-parameter series $y(a, x)$ given by (130), we have the following relation:

$$y(a^n, y_n(x)) = y_n(y(a, x)) = a^n \cdot x^n + 744 \cdot n \cdot a^n \cdot x^{n+1} + \dots \quad (187)$$

5.5. Composition in general

The one-parameter series (186) can be written

$$y_n^{(a)}(x) = \tilde{X}(a \cdot \tilde{Q}(x)^n). \quad (188)$$

We have the following composition:

$$\begin{aligned} y_n^{(a)}(y_m^{(b)}(x)) &= \tilde{X}(a \cdot \tilde{Q}(\tilde{X}(b \cdot \tilde{Q}(x)^m))^n) = \tilde{X}(a \cdot (b \cdot \tilde{Q}(x)^m)^n) \\ &= \tilde{X}(a \cdot b^n \cdot \tilde{Q}(x)^{mn}) = y_{mn}^{(ab^n)}(x). \end{aligned} \quad (189)$$

Note that the condition to have series solutions of the Schwarzian equation of the form $y_n^{(a)}(x) = a \cdot x^n + \dots$, with $n \geq 2$, amounts to having [30, 31] $W(x)$ of the form $W(x) = -1/2x^2 + \dots$ which is satisfied when $F(x) = \alpha \cdot x + \dots$, or $\tilde{Q}(x) = \rho \cdot x^{1/\alpha} + \dots$

6. The one-parameter series (4) seen as a ϵ -expansion.

In the $a \rightarrow 1$ limit, let us denote $\epsilon = a - 1$. The one-parameter series $y(x) = y(a, x)$, given by (130), can, thus, be seen as an ϵ -expansion:

$$y(a, x) = x + \sum_{n=1}^{\infty} \epsilon^n \cdot B_n(x), \quad (190)$$

where $B_1(x) = F(x)$, with $F(x)$ given by (168), and where $B_2(x)$ reads (see equation (115) in [30]):

$$B_2(x) = \frac{1}{2} \cdot F(x) \cdot \left(\frac{dB_1(x)}{dx} - 1 \right). \quad (191)$$

Assuming that (190) is solution of the Schwarzian condition (174) (with $W(x)$ given by (170)), we actually obtained the next $B_n(x)$'s:

$$\begin{aligned} B_3(x) &= \frac{1}{3} \cdot F(x) \cdot \left(\frac{dB_2(x)}{dx} - \frac{dB_1(x)}{dx} + 1 \right), \\ B_4(x) &= \frac{1}{4} \cdot F(x) \cdot \left(\frac{dB_3(x)}{dx} - \frac{dB_2(x)}{dx} + \frac{dB_1(x)}{dx} - 1 \right), \\ B_5(x) &= \frac{1}{5} \cdot F(x) \cdot \left(\frac{dB_4(x)}{dx} - \frac{dB_3(x)}{dx} + \frac{dB_2(x)}{dx} - \frac{dB_1(x)}{dx} + 1 \right), \\ B_6(x) &= \frac{1}{6} \cdot F(x) \cdot \left(\frac{dB_5(x)}{dx} - \frac{dB_4(x)}{dx} + \frac{dB_3(x)}{dx} - \frac{dB_2(x)}{dx} + \frac{dB_1(x)}{dx} - 1 \right), \quad \dots \end{aligned} \quad (192)$$

More generally, one easily discovers the recursion

$$(n+1) \cdot B_{n+1} + n \cdot B_n = F(x) \cdot \frac{dB_n(x)}{dx}, \quad (193)$$

which yields on the series (190)

$$\sum_n (n+1) \cdot B_{n+1} \cdot \epsilon^n + \sum_n n \cdot B_n \cdot \epsilon^n = F(x) \cdot \left(\sum_n \frac{dB_n(x)}{dx} \cdot \epsilon^n \right), \quad (194)$$

or

$$\frac{\partial \sum_n B_{n+1} \cdot \epsilon^{n+1}}{\partial \epsilon} + \epsilon \cdot \frac{\partial \sum_n B_n \cdot \epsilon^n}{\partial \epsilon} = F(x) \cdot \left(\frac{\partial \sum_n B_n(x) \cdot \epsilon^n}{\partial x} \right), \quad (195)$$

yielding finally

$$(1 + \epsilon) \cdot \frac{\partial y(a, x)}{\partial \epsilon} = F(x) \cdot \frac{\partial y(a, x)}{\partial x}, \quad (196)$$

namely:

$$a \cdot \frac{\partial y(a, x)}{\partial a} = F(x) \cdot \frac{\partial y(a, x)}{\partial x}. \quad (197)$$

Note that $y(a, x)$ is also solution of:

$$F(y(a, x)) = F(x) \cdot \frac{\partial y(a, x)}{\partial x}. \quad (198)$$

Recalling some relation on the nome q (see equation (33) in [30]):

$$\frac{q'}{q} = \frac{1}{F(x)} \quad \text{or:} \quad q \cdot \frac{d}{dq} = F(x) \cdot \frac{d}{dx}, \quad (199)$$

we see that relation (197) also reads more simply:

$$a \cdot \frac{\partial y(a, x)}{\partial a} = q \cdot \frac{\partial y(a, x)}{\partial q}. \quad (200)$$

which is reminiscent of the fact that changing $x \rightarrow y(a, x)$ just amounts, on the nome, to changing $q \rightarrow a \cdot q$. Equation (197) means that $y(a, x)$ is a function of

$$\int \left(\frac{da}{a} + \frac{dx}{F(x)} \right) = \ln(a) + \int \left(\frac{dx}{F(x)} \right), \quad (201)$$

or, recalling (171), a function of:

$$\exp \left(\int \left(\frac{da}{a} + \frac{dx}{F(x)} \right) \right) = a \cdot \tilde{Q}(x). \quad (202)$$

This is actually the case since $y(a, x)$ is nothing but $\tilde{X}(a \cdot \tilde{Q}(x))$ (see (133)).

Remark 6.1: Do note that the previous calculations *are still valid* when $F(x)$ is *not given by* (168). One can verify, for *any function* $F(x)$, that the ϵ -expansion (190) with coefficients B_n given by (191), (192), (193), *is actually solution of the Schwarzian relation* (174), with $W(x)$ given by:

$$W(x) = \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)} \right)^2. \quad (203)$$

7. Generalization of $W(x)$ in the Schwarzian equation: adding an extra parameter α .

For a given function $F(x)$ let us consider the relation

$$F(y(x)) = F(x) \cdot \frac{dy(x)}{dx}, \quad (204)$$

which corresponds to:

$$\frac{dy}{F(y)} = \frac{dx}{F(x)} = \frac{dq}{q}. \quad (205)$$

From (204), namely $F(y) = F(x) \cdot y'$, one gets

$$F'(y) \cdot y' = F'(x) \cdot y' + F(x) \cdot y'', \quad (206)$$

or

$$\frac{F'(y)}{F(y)} \cdot y' = \frac{F'(x)}{F(x)} + \frac{y''}{y'}. \quad (207)$$

or, more generally, using (204) in order to introduce an extra parameter α :

$$\left(\frac{F'(y)}{F(y)} + \frac{\alpha}{F(y)} \right) \cdot y' = \left(\frac{F'(x)}{F(x)} + \frac{\alpha}{F(x)} \right) + \frac{y''}{y'}. \quad (208)$$

Let us introduce

$$A_R(x) = \frac{F'(x)}{F(x)} + \frac{\alpha}{F(x)}, \quad (209)$$

we see that (208) can be written

$$A_R(x) - A_R(y) \cdot y' + \frac{y''}{y'} = 0. \quad (210)$$

which is (32) of section (1.6). From (206), that we rewrite

$$F'(y) = F'(x) + F(x) \cdot \frac{y''}{y'}, \quad (211)$$

one gets

$$F''(y) \cdot y' = F''(x) + F'(x) \cdot \frac{y''}{y'} + F(x) \cdot \left(\frac{y''}{y'}\right)', \quad (212)$$

or, using (204), written $F(x) = F(y)/y'$:

$$\frac{F''(y)}{F(y)} \cdot y'^2 = \frac{F''(x)}{F(x)} + \frac{F'(x)}{F(x)} \cdot \frac{y''}{y'} + \left(\frac{y''}{y'}\right)'. \quad (213)$$

Taking the square of (207) one gets (up to a factor 2):

$$\frac{1}{2} \cdot \left(\frac{F'(y)}{F(y)}\right)^2 \cdot y'^2 = \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)}\right)^2 + \frac{1}{2} \cdot \left(\frac{y''}{y'}\right)^2 + \frac{F'(x)}{F(x)} \cdot \frac{y''}{y'}. \quad (214)$$

From (213) and (214) we deduce:

$$\left(\frac{F''(y)}{F(y)} - \frac{1}{2} \cdot \left(\frac{F'(y)}{F(y)}\right)^2\right) \cdot y'^2 = \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)}\right)^2 + \left(\frac{y''}{y'}\right)' - \frac{1}{2} \cdot \left(\frac{y''}{y'}\right)^2,$$

or, recalling the Schwarzian derivative,

$$\frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)}\right)^2 - \left(\frac{F''(y)}{F(y)} - \frac{1}{2} \cdot \left(\frac{F'(y)}{F(y)}\right)^2\right) \cdot y'^2 + \{y(x), x\} = 0,$$

or, more generally, using (204), which allows to introduce an extra parameter α

$$\begin{aligned} & \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)}\right)^2 + \frac{1}{2} \cdot \frac{\alpha^2}{F(x)^2} \\ & - \left(\frac{F''(y)}{F(y)} - \frac{1}{2} \cdot \left(\frac{F'(y)}{F(y)}\right)^2 + \frac{1}{2} \cdot \frac{\alpha^2}{F(y)^2}\right) \cdot y'^2 + \{y(x), x\} = 0. \end{aligned} \quad (215)$$

Note that (215) is actually of the form

$$W(x) - W(y(x)) \cdot y'(x)^2 + \{y(x), x\} = 0, \quad (216)$$

where $(A_R$ given by (209)):

$$W(x) = \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)}\right)^2 + \frac{1}{2} \cdot \frac{\alpha^2}{F(x)^2} = A'_R(x) + \frac{A_R(x)^2}{2}. \quad (217)$$

Remark 7.1: Note that these calculations *also work* with

$$\mu \cdot F(y(x)) = F(x) \cdot \frac{dy(x)}{dx}, \quad (218)$$

which corresponds to (49), (59), (77), (85).

8. An “academical” Schwarzian equation: $W(x)$ is no longer a rational function

Recalling

$$F(x) = x \cdot (1 - 1728 \cdot x)^{1/2} \cdot {}_2F_1\left(\left[\frac{1}{12}, \frac{5}{12}\right], [1], 1728 \cdot x\right)^2, \quad (219)$$

the one-parameter series $y(x) = y(a, x)$, given by (130), is, for any value of α , solution of the rank-two equation

$$A_R(x) - A_R(y(x)) \cdot y'(x) + \frac{y''(x)}{y'(x)} = 0, \quad (220)$$

with

$$A_R(x) = \frac{F'(x)}{F(x)} + \frac{\alpha}{F(x)}, \quad (221)$$

but is also solution of the Schwarzian condition

$$W(x) - W(y(x)) \cdot y'(x)^2 + \{y(x), x\} = 0, \quad (222)$$

where:

$$\begin{aligned} W(x) &= \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)} \right)^2 + \frac{1}{2} \cdot \frac{\alpha^2}{F(x)} = A'_R(x) + \frac{A_R(x)^2}{2} \\ &= -\frac{1}{2} \cdot \frac{1 - 1968x + 2654208x^2}{x^2 \cdot (1 - 1728x)^2} + \frac{1}{2} \cdot \frac{\alpha^2}{F(x)}. \end{aligned} \quad (223)$$

For *generic* values of α , the solution-series of the form $a \cdot x + \dots$, of the rank-two equation (220) with (221), as well as the Schwarzian equation (222), with $W(x)$ given by (223), is just the one-parameter series $y(x) = y(a, x)$, given by (130). However, for a selected set of values of α , namely (non-zero) *integer values*, the solution-series of the form $a \cdot x + \dots$, becomes a *two-parameters series*. For instance, for $\alpha = \pm 1$, the extra parameter occurs with the coefficient of x^2 , for $\alpha = \pm 2$ the extra parameter occurs with the coefficient of x^3 , ... and, more generally, for $\alpha = \pm N$ the extra parameter occurs with the coefficient of x^{N+1} . Let us display the $\alpha = 1$ case in detail.

8.1. The $\alpha = 1$ case: two-parameters series

Let us consider the case $\alpha = 1$ in the Schwarzian equation (222) with $W(x)$ given by (223), or in the rank-two relation (220) with (221).

The *two-parameter series*

$$\begin{aligned} y(a, b, x) &= a \cdot x + \left(1728 \cdot b - 744 \cdot a \cdot (a - 1) \right) \cdot x^2 \\ &\quad + \left(2985984 \cdot a \cdot b^2 - 2571264 \cdot a \cdot (a - 1) \cdot b \right. \\ &\quad \left. + 36 \cdot a \cdot (a - 1) \cdot (9907a - 20845) \right) \cdot x^3 \\ &\quad + \left(5159780352 \cdot a \cdot b^3 - 6664716288 \cdot a \cdot (a - 1) \cdot b^2 \right. \\ &\quad \left. + 186624 \cdot (9907a^2 - 30752a + 19022) \cdot a \cdot b \right. \\ &\quad \left. - 32 \cdot a \cdot (a - 1) \cdot (4386286a^2 - 20490191a + 27274051) \right) \cdot x^4 + \dots \end{aligned} \quad (224)$$

is *actually solution of the Schwarzian equation (222)* with $W(x)$ given by (223), or of the rank-two relation (220) with (221), for $\alpha = 1$. Note that the *two-parameter series (224)* is also solution[†] of

$$a \cdot \frac{\partial y(a, b, x)}{\partial a} + b \cdot \frac{\partial y(a, b, x)}{\partial b} = F(y(a, b, x)), \quad (225)$$

with $F(x)$ given by (219). We have the following composition rules for the *two-parameter series (224)*:

$$y(a', b', y(a, b, x)) = y(a a', a^2 b' + a' b, x). \quad (226)$$

[†] However it is *not* solution of $F(x) \cdot y' = F(y)$ or $F(x) \cdot y' = a \frac{\partial y}{\partial a}$.

Let us introduce an *alternative parametrization* of the two-parameter series (224), changing b into $a b$, in (224):

$$\begin{aligned} Y(a, b, x) = & a \cdot x + \left(1728 \cdot a b - 744 \cdot a \cdot (a-1)\right) \cdot x^2 \\ & + \left(2985984 \cdot a^3 \cdot b^2 - 2571264 \cdot a^2 \cdot (a-1) \cdot b \right. \\ & \left. + 36 \cdot a \cdot (a-1) \cdot (9907 a - 20845)\right) \cdot x^3 + \dots \end{aligned} \quad (227)$$

We have the following composition rules for the *two-parameter* series (227)

$$Y(a', b', Y(a, b, x)) = Y(a a', a b' + b, x). \quad (228)$$

The series (227) is, now, solution of:

$$a \cdot \frac{\partial Y(a, b, x)}{\partial a} = F(Y(a, b, x)), \quad (229)$$

with $F(x)$ given by (219). Let us introduce the $a \rightarrow 0$ limit:

$$\begin{aligned} Q_b(x) = & \lim_{a \rightarrow 0} \frac{Y(a, b, x)}{a} \\ = & x + (744 + 1728 b) \cdot x^2 + (750420 + 2571264 b + 2985984 b^2) \cdot x^3 \\ & + (872769632 + 3549961728 b + 6664716288 b^2 + 5159780352 b^3) \cdot x^4 \\ & + (1102652742882 + 4945819779072 b + 11680775258112 b^2 \\ & + 15355506327552 b^3 + 8916100448256 b^4) \cdot x^5 \\ & + (1470561136292880 + 7027977959274240 b + 19050621395927040 b^2 \\ & + 32624754548539392 b^3 + 33167893667512320 b^4 + 15407021574586368 b^5) \cdot x^6 \\ & + \dots \end{aligned} \quad (230)$$

In the $b \rightarrow 0$ limit, this series (230) reduces to the nome series (135) or (136).

In the $a \rightarrow \infty$ limit one gets:

$$\begin{aligned} X_b(x) = & \lim_{a \rightarrow \infty} Y\left(a, b, \frac{x}{a}\right) = x - 744 x^2 + 356652 x^3 - 140361152 x^4 \\ & + 49336682190 x^5 - 16114625669088 x^6 + 4999042477430456 x^7 \\ & - 1492669384085015040 x^8 + 432762759484818142437 x^9 + \dots \end{aligned} \quad (231)$$

This series (231) is nothing but (134) or (137), and, thus, *does not depend* on the second parameter b .

One actually finds that the two parameter series (227) is *nothing but*:

$$Y(a, b, x) = X_b(a \cdot Q_b(x)). \quad (232)$$

From (232) we can also deduce that (229), is, in fact, nothing but equation:

$$a \cdot \frac{\partial X_b(a \cdot x)}{\partial a} = F(X_b(a \cdot x)). \quad (233)$$

Furthermore, since $a \cdot \frac{\partial X_b(a \cdot x)}{\partial a} = x \cdot \frac{\partial X_b(a \cdot x)}{\partial x}$, relation (233) also gives:

$$x \cdot \frac{\partial X_b(a \cdot x)}{\partial x} = F(X_b(a \cdot x)). \quad (234)$$

In contrast with the $b = 0$ case, the two functions, Q_b and X_b , given by the two limits (230), (231), are *not compositional inverse*. In the $a \rightarrow 1$ limit, the decomposition (232) becomes:

$$\begin{aligned} Y(1, b, x) &= X_b(Q_b(x)) \\ &= x + 1728 \cdot b \cdot x^2 + 2985984 \cdot b^2 \cdot x^3 + 186624 \cdot (27648b^2 - 1823) \cdot b \cdot x^4 \\ &\quad + 110592 \cdot (80621568b^3 - 15947604b - 5249233) \cdot b \cdot x^5 + \dots \end{aligned} \quad (235)$$

The series (235) is a one-parameter family of commuting series:

$$Y(1, b, Y(1, b', x)) = Y(1, b', Y(1, b, x)) = Y(1, b + b', x). \quad (236)$$

In particular the compositional inverse of $Y(1, b, x)$ is $Y(1, -b, x)$:

$$Y(1, b, Y(1, -b, x)) = Y(1, -b, Y(1, b, x)) = x. \quad (237)$$

Note that:

$$Q_b(X_b(x)) = \frac{x}{1 - 1728 \cdot b \cdot x} = x + 1728 \cdot b \cdot x^2 + \dots \quad (238)$$

From (238) we deduce an alternative expression for $Q_b(x)$ in terms of the nome (23) (i.e. the compositional inverse of (231), or, equivalently $Q_b(x)$ for $b = 0$):

$$Q_b(x) = \frac{Q_0(x)}{1 - 1728 \cdot b \cdot Q_0(x)}. \quad (239)$$

Conversely, the nome (23), *which does not depend on the parameter b* , can be simply expressed in terms of the series (230):

$$Q_0(x) = \frac{Q_b(x)}{1 + 1728 \cdot b \cdot Q_b(x)}. \quad (240)$$

Note that the composition rule relation (228) can, now, be seen as a straightforward consequence of relation (239). From relation (239) one can see that the radius of convergence of the series (230) corresponds, for small enough values of the additional parameter b , to the singularity of $Q_0(x)$, (i.e. $R = 1/1728$), and for large enough values of the parameter b , to the singularity $Q_0(x) = 1/1728/b$, namely:

$$x = X_b\left(\frac{1}{1728b}\right) = \tilde{X}\left(\frac{1}{1728b}\right). \quad (241)$$

Remark 8.1: Do note that, in contrast with the $\alpha = 0$ case, *there is no* solution-series of the form $a \cdot x^2 + \dots$ or, more generally, of the form $a \cdot x^N + \dots$ with $N \neq 1$, of the Schwarzian equation (222), when $W(x)$ is given by (223). This corresponds to the fact that, when $\alpha \neq 0$, $W(x)$ is no longer of the form $W(x) = -1/2/x^2 + \dots$ (see [30, 31]).

9. Polynomial examples for $F(x)$.

Modular correspondences, modular curves, correspond to a (transcendental) function $F(x)$ associated to elliptic functions like (168), (219).

Appendix B provides a (non globally bounded) Heun function example showing that the previous results and calculations also work, *mutatis mutandis* with Shimura curves [75] (and their associated automorphic forms [10]).

Let us now recall the general results of section (6), which describes the *one-parameter* solution-series (190) of the Schwarzian equation (174), and also the partial differential equations (197), (198), and the fact that these equations *are actually valid for any function* $F(x)$.

Let us consider, here, one-parameter functions $y(a, x)$, corresponding to miscellaneous *polynomial* examples of functions $F(x)$, that are, thus, far from being associated with the previous “classical” modular forms [14, 15, 19] and hypergeometric/elliptic functions [45, 46], or even Shimura curves/automorphic forms examples, possibly with Heun functions [83] (see Appendix B). Even more simple polynomial examples are given in Appendix C.

From the general results of the previous section (6) we will, thus, get a set of miscellaneous examples. All the corresponding one-parameter series, below, will verify the composition rule:

$$y\left(a, y(a', x)\right) = y(a a', x). \quad (242)$$

All these one-parameter series will also verify:

$$F\left(y(a, x)\right) = a \cdot \frac{\partial y(a, x)}{\partial a} = F(x) \cdot \frac{\partial y(a, x)}{\partial x}. \quad (243)$$

One will also consider a polynomial that will be the truncation of the hypergeometric function (168). One will, then, get a one-parameter solution-series, very similar[‡] to (130), which also verifies the composition rule (132), but does not correspond to globally bounded series [43].

9.1. A first simple polynomial example for $F(x)$

Let us first consider the following polynomial expression for $F(x)$:

$$F(x) = x \cdot (1 - 2x) \cdot (1 - 3x). \quad (244)$$

One deduces, from (169), (203), the following rational expression for $W(x)$:

$$W(x) = -\frac{1}{2} \cdot \frac{1 - 36x^2 + 120x^3 - 108x^4}{x^2 \cdot (1 - 2x)^2 \cdot (1 - 3x)^2}. \quad (245)$$

The Schwarzian condition

$$W(x) - W(y(x)) \cdot y'(x)^2 + \{y(x), x\} = 0, \quad (246)$$

has the following one-parameter solution

$$\begin{aligned} y(a, x) = & a \cdot x - 5 \cdot a \cdot (a - 1) \cdot x^2 + 2 \cdot a \cdot (14a - 11)(a - 1) \cdot x^3 \\ & - 15 \cdot a \cdot (a - 1)^2 (11a - 6) \cdot x^4 + a \cdot (a - 1)^2 \cdot (1001a^2 - 1298a + 351) \cdot x^5 \\ & - 7 \cdot a \cdot (a - 1)^3 \cdot (884a^2 - 923a + 189) \cdot x^6 \\ & + 60 \cdot a \cdot (a - 1)^3 \cdot (646a^3 - 1156a^2 + 600a - 81) \cdot x^7 \\ & - 3 \cdot a \cdot (a - 1)^4 \cdot (81719a^3 - 125324a^2 + 54162a - 5832) \cdot x^8 + \dots \end{aligned} \quad (247)$$

as well as modular equation-like series (with no parameter) like

$$\begin{aligned} y_2 = & x^2 + 10x^3 + 64x^4 + 300x^5 + 924x^6 + 56x^7 - 24140x^8 \\ & - 209856x^9 - 1158600x^{10} + \dots \end{aligned} \quad (248)$$

[‡] The three first terms are the same.

or:

$$y_3 = x^3 + 15x^4 + 141x^5 + 1050x^6 + 6705x^7 + 37854x^8 + 189603x^9 + 820584x^{10} + 2777004x^{11} + 4024890x^{12} + \dots \quad (249)$$

More generally, for the solution series of the form $x \rightarrow x^p + \dots$ we have:

$$p \cdot \frac{dx}{F(x)} = \frac{dy}{F(y)}, \quad (250)$$

From (244) we get:

$$\exp\left(\int \frac{dx}{F(x)}\right) = \mu \cdot \frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}. \quad (251)$$

From (250), and (251), we get the *algebraic curve*:

$$\rho \cdot \left(\frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}\right)^p = \frac{y \cdot (1 - 2y)^2}{(1 - 3y)^3}. \quad (252)$$

One finds that y_2 , given by (248), is actually solution of the algebraic condition (252) for $p = 2$ and $\rho = 1$, namely:

$$\left(\frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}\right)^2 = \frac{y \cdot (1 - 2y)^2}{(1 - 3y)^3}. \quad (253)$$

One also finds that y_3 , given by (249), is actually solution of the algebraic condition (252), for $p = 3$ and $\rho = 1$, namely:

$$\left(\frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}\right)^3 = \frac{y \cdot (1 - 2y)^2}{(1 - 3y)^3}. \quad (254)$$

On the other hand one finds that the one-parameter series $y(a, x)$, given by (247), is actually solution of the algebraic condition (252) for $p = 1$ and $\rho = a$, namely:

$$a \cdot \left(\frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}\right) = \frac{y \cdot (1 - 2y)^2}{(1 - 3y)^3}. \quad (255)$$

Remark 9.1: Note, from (255), that the one-parameter series $y(a, x)$, given by (247), is actually an *algebraic series* for *any value* of the parameter a (and not only N -th root of unity). The algebraic equations (253) and (254), and their corresponding algebraic series solutions (248) and (249), could be seen to be the “equivalent” of the modular equations (43) and (53), and their corresponding algebraic series solutions (45) and (55). However, one should note that the *modular equations* (43) and (53) are $x \leftrightarrow y$ symmetric, and, consequently, the modular equation (43) represents $q \rightarrow q^2$ and $q \rightarrow q^{1/2}$ in the same time (see series (45) but also (46)). Similarly the modular equation (53) represents $q \rightarrow q^3$ and $q \rightarrow q^{1/3}$ in the same time (see series (55) and also (56)). In contrast (253) and (254) break the $x \leftrightarrow y$ symmetry. Therefore, the “equivalent” of the modular equation (43) is rather

$$\left(\left(\frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}\right)^2 - \frac{y \cdot (1 - 2y)^2}{(1 - 3y)^3}\right) \cdot \left(\left(\frac{y \cdot (1 - 2y)^2}{(1 - 3y)^3}\right)^2 - \frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}\right) = 0,$$

when the equivalent of the modular equation (53) is rather:

$$\left(\left(\frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}\right)^3 - \frac{y \cdot (1 - 2y)^2}{(1 - 3y)^3}\right) \cdot \left(\left(\frac{y \cdot (1 - 2y)^2}{(1 - 3y)^3}\right)^3 - \frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}\right) = 0.$$

The series (247) verifies the composition rule:

$$y(a, y(a', x)) = y(a a', x). \quad (256)$$

The series (247) also verifies the relations:

$$F\left(y(a, x)\right) = F(x) \cdot \frac{\partial y(a, x)}{\partial x} = a \cdot \frac{\partial y(a, x)}{\partial a}. \quad (257)$$

Let us introduce the two limits:

$$\begin{aligned} \tilde{Q}(x) = \lim_{a \rightarrow 0} \frac{y(a, x)}{a} = & x + 5x^2 + 22x^3 + 90x^4 + 351x^5 + 1323x^6 \\ & + 4860x^7 + 17496x^8 + 61965x^9 + 216513x^{10} + \dots \end{aligned} \quad (258)$$

$$\begin{aligned} \tilde{X}(x) = \lim_{a \rightarrow \infty} y\left(a, \frac{x}{a}\right) = & x - 5x^2 + 28x^3 - 165x^4 + 1001x^5 - 6188x^6 \\ & + 38760x^7 - 245157x^8 + 1562275x^9 - 10015005x^{10} + \dots \end{aligned} \quad (259)$$

One can verify that these two series $\tilde{Q}(x)$ and $\tilde{X}(x)$ are *compositional inverse*. The radius of convergence of the “nome-like” series $\tilde{Q}(x)$, given by (258), is $R = 1/3$. The radius of convergence of the series $\tilde{X}(x)$, given by (259), is $R = 4/27$.

These two series, with *integer* coefficients, are solutions of the two Schwarzian equations

$$\{\tilde{Q}(x), x\} + \frac{1}{2 \cdot \tilde{Q}(x)^2} \cdot \left(\frac{d\tilde{Q}(x)}{dx}\right)^2 + W(x) = 0, \quad (260)$$

and

$$\{\tilde{X}(x), x\} - \frac{1}{2 \cdot x^2} - W(\tilde{X}(x)) \cdot \left(\frac{d\tilde{X}(x)}{dx}\right)^2 = 0, \quad (261)$$

where $W(x)$ is given by (245). In fact using the *explicit algebraic form* of $y(a, x)$, given by (255), one can find a closed exact expression for the “nome-like” series $\tilde{Q}(x)$, namely:

$$\tilde{Q}(x) = \frac{x \cdot (1 - 2x)^2}{(1 - 3x)^3}, \quad (262)$$

in agreement with series (258). Relation (255) is nothing but:

$$a \cdot \tilde{Q}(x) = \tilde{Q}\left(y(a, x)\right). \quad (263)$$

The one-parameter series (247) can thus be written:

$$y(a, x) = \tilde{X}\left(a \cdot \tilde{Q}(x)\right). \quad (264)$$

From (264), and from the fact that $y(a, x) = x$ for $a = 1$, one can deduce that $\tilde{X}(x)$ *must be the compositional inverse* of the “nome-like” series (262). Note that $\tilde{X}(x)$ is an algebraic function. It is solution of the polynomial equation:

$$(27x + 4) \cdot \tilde{X}(x)^3 - (27x + 4) \cdot \tilde{X}(x)^2 + (9x + 1) \cdot \tilde{X}(x) - x = 0, \quad (265)$$

in agreement with the $R = 4/27$ radius of convergence of the series $\tilde{X}(x)$.

9.2. Another simple polynomial example for $F(x)$

Let us now consider the polynomial

$$F(x) = x \cdot (1 - 373 \cdot x) \cdot (1 - 371 \cdot x) = x - 744x^2 + 138383x^3, \quad (266)$$

which has the *same first two terms* as the series expansion of the hypergeometric function (168). The function $W(x)$ in the Schwarzian equation is given by (203):

$$W(x) = -\frac{1}{2} \cdot \frac{1 - 830298x^2 + 411827808x^3 - 57449564067x^4}{x^2 \cdot (1 - 373x)^2 \cdot (1 - 371x)^2}. \quad (267)$$

A solution of the Schwarzian equation, with $W(x)$ given by (267), reads:

$$\begin{aligned} y(a, x) = & a \cdot x - 744 \cdot a \cdot (a-1) \cdot x^2 + \frac{1}{2} \cdot a \cdot (1245455a - 968689) \cdot (a-1) \cdot x^3 \\ & - 620 \cdot a \cdot (885656a - 470507) \cdot (a-1)^2 \cdot x^4 + \dots \end{aligned} \quad (268)$$

The functional relation $F(y) = F(x) \cdot y'(x)$, gives $dy/F(y) = dx/F(x)$, and thus

$$\begin{aligned} \mu \cdot \exp\left(\int \frac{dy}{F(y)}\right) &= \tilde{Q}(x) = \exp\left(\int \frac{dx}{F(x)}\right) = x \cdot \frac{(1 - 371x)^{371/2}}{(1 - 373x)^{373/2}} \\ &= x + 744x^2 + \frac{968689}{2}x^3 + 291714340x^4 + \dots \end{aligned} \quad (269)$$

We actually have the relation:

$$a \cdot \tilde{Q}(x) = \tilde{Q}(y(a, x)). \quad (270)$$

In that case, since $\tilde{Q}(x)$ is an *algebraic function*, we see that the one-parameter series $y(a, x)$, given by (268), *is actually an algebraic series for any value of the parameter a* . The series $y = y(a, x)$ is actually solution of:

$$a^2 \cdot x^2 \cdot \frac{(1 - 371x)^{371}}{(1 - 373x)^{373}} - y^2 \cdot \frac{(1 - 371y)^{371}}{(1 - 373y)^{373}} = 0. \quad (271)$$

Taking into account the large degree in x or y of the polynomial condition (271), one should note that it can actually be quite difficult to get this polynomial equation from ¶ a large series (268). The *compositional inverse* of $\tilde{Q}(x)$ is:

$$\begin{aligned} \tilde{X}(x) = & x - 744x^2 + \frac{1245455}{2} \cdot x^3 - 549106720x^4 + \frac{3989599188003}{8} \cdot x^5 \\ & - 461623555588416x^6 + \frac{6928370820171415659}{16} \cdot x^7 \\ & - 410201463628637176320x^8 + \dots \end{aligned} \quad (272)$$

This is an *algebraic series* $y = \tilde{X}(x)$, solution of:

$$x^2 \cdot (1 - 373 \cdot y)^{373} - y^2 \cdot (1 - 371 \cdot y)^{371} = 0. \quad (273)$$

Note that, even with a very large series (272), it is also quite hard, because of the high degree in y of (273), to find the algebraic expression (273) even if it is really simple.

9.2.1. Two-parameter family. Following the calculations displayed in subsection (8.1), let us generalize $W(x)$ given by (267), to the form (223):

$$W(x) = \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)}\right)^2 + \frac{1}{2} \cdot \frac{\alpha^2}{F(x)} \quad (274)$$

For $\alpha = 1$ with $F(x)$ given by (266), $W(x)$ reads:

$$W(x) = \frac{415149}{2} \cdot \frac{138383x^2 - 992x + 2}{x^2 \cdot (1 - 373x)^2 \cdot (1 - 371x)^2}. \quad (275)$$

¶ Using, for instance, the command `seriestoalgeq` of `gfun` of Bruno Salvy.

The *two-parameter* series (generalizing (268))

$$\begin{aligned}
y(a, b, x) = & a \cdot x + \left(1728 \cdot b - 744 \cdot a \cdot (a-1)\right) \cdot x^2 \\
& + \left(2985984 \cdot \frac{b^2}{a} - 2571264 \cdot (a-1) \cdot b \right. \\
& \quad \left. + \frac{1}{2} \cdot a \cdot (1245455 a - 968689) \cdot (a-1)\right) \cdot x^3 \\
& + \left(5159780352 \cdot \frac{b^3}{a^2} - 6664716288 \cdot (a-1) \cdot \frac{b^2}{a} \right. \\
& \quad + 864 \cdot (3736365 a^2 - 6642432 a + 3044450) \cdot b \\
& \quad \left. - 620 \cdot a \cdot (885656 a - 470507) \cdot (a-1)^2\right) \cdot x^4 + \dots
\end{aligned} \tag{276}$$

is *actually* solution of the *Schwarzian equation* (222) with $W(x)$ given by (275). Note that, again, the *two-parameter* series (276) is also solution of

$$a \cdot \frac{\partial y(a, b, x)}{\partial a} + b \cdot \frac{\partial y(a, b, x)}{\partial b} = F(y(a, b, x)), \tag{277}$$

with $F(x)$ given by (266). We have, again, the *two-parameters composition rules* (226):

$$y(a', b', y(a, b, x)) = y(a a', a^2 b' + a' b, x). \tag{278}$$

similarly to subsection (8.1), let us introduce an *alternative parametrization* of the two-parameter series (276), changing b into $a b$, in (276):

$$\begin{aligned}
Y(a, b, x) = & a \cdot x + \left(1728 \cdot a b - 744 \cdot a \cdot (a-1)\right) \cdot x^2 \\
& + \frac{1}{2} \cdot \left(5971968 \cdot a \cdot b^2 - 5142528 \cdot (a-1) \cdot a \cdot b \right. \\
& \quad \left. + a \cdot (1245455 a - 968689) \cdot (a-1)\right) \cdot x^3 + \dots
\end{aligned} \tag{279}$$

Again, we have the following composition rules for the *two-parameter* series (279)

$$Y(a', b', Y(a, b, x)) = Y(a a', a b' + b, x), \tag{280}$$

The series (279) is, now, solution of:

$$a \cdot \frac{\partial Y(a, b, x)}{\partial a} = F(Y(a, b, x)), \tag{281}$$

with $F(x)$ given by (266).

9.3. Truncation of the hypergeometric function $F(x)$.

The hypergeometric function $F(x)$ given by (168), expands as $x - 744 x^2 - 393768 x^3 + \dots$ Let us consider a simple truncation of this hypergeometric function:

$$F(x) = x - 744 x^2 - 393768 x^3. \tag{282}$$

From (203) this gives:

$$W(x) = -\frac{1}{2} \cdot \frac{1 + 2362608 x^2 - 1171853568 x^3 - 465159713472 x^4}{x^2 \cdot (1 - 744 x - 393768 x^2)^2}. \tag{283}$$

The Schwarzian equation (174) with the previous $W(x)$, namely (283), has the following *one-parameter* solution-series:

$$\begin{aligned} y(a, x) = & a \cdot x - 744 \cdot a \cdot (a-1) \cdot x^2 + 36 \cdot a \cdot (a-1) \cdot (9907a - 20845) \cdot x^3 \\ & - 80352 \cdot a \cdot (a-1)^2 \cdot (264a - 9379) \cdot x^4 \\ & - 648 \cdot a \cdot (a-1)^2 \cdot (250310357a^2 + 598043050a - 1207272939) \cdot x^5 \\ & + \frac{482112}{5} \cdot a \cdot (a-1)^3 \cdot (1944308192a^2 - 424834349a - 8498464743) \cdot x^6 + \dots \end{aligned} \quad (284)$$

This one-parameter series (284) is quite similar[†] to the one-parameter series (130). The series (284) actually verifies the composition rule:

$$y(a, y(a', x)) = y(a a', x). \quad (285)$$

Let us introduce the two limits

$$\begin{aligned} \tilde{Q}(x) = \lim_{a \rightarrow 0} \frac{y(a, x)}{a} = & x + 744x^2 + 750420x^3 + 753621408x^4 \\ & + 782312864472x^5 + \frac{4097211834177216}{5}x^6 + \frac{4331866321367059104}{5}x^7 + \dots \end{aligned} \quad (286)$$

and:

$$\begin{aligned} \tilde{X}(x) = \lim_{a \rightarrow \infty} y\left(a, \frac{x}{a}\right) = & x - 744x^2 + 356652x^3 - 21212928x^4 \\ & - 162201111336x^5 + \frac{937374311061504}{5}x^6 - \frac{563689525139743392}{5}x^7 + \dots \end{aligned} \quad (287)$$

One verifies that the one-parameter series (284) *is actually of the form*:

$$y(a, x) = \tilde{X}\left(a \cdot \tilde{Q}(x)\right). \quad (288)$$

Again, from (288) and from the fact that $y(a, x) = x$ for $a = 1$, we see that the series (287) is actually the *compositional inverse* of the “nome-like” series (286):

$$y(1, x) = x = \tilde{X}\left(\tilde{Q}(x)\right). \quad (289)$$

The *one-parameter* series (284) is also solution of

$$a \cdot \frac{\partial y(a, x)}{\partial a} = F(y(a, x)) = F(x) \cdot \frac{\partial y(a, x)}{\partial x}, \quad (290)$$

and one can verify that:

$$\tilde{Q}\left(y(a, x)\right) = a \cdot \tilde{Q}(x). \quad (291)$$

Conversely, from (291), we get, recalling (205)

$$\frac{dy}{F(y(a, x))} = \frac{dx}{F(x)} + \frac{da}{a}, \quad (292)$$

which gives for a fixed

$$F(y(a, x)) = F(x) \cdot \frac{\partial y(a, x)}{\partial x}, \quad (293)$$

and for x fixed:

$$a \cdot \frac{\partial y(a, x)}{\partial a} = F(y(a, x)). \quad (294)$$

[†] The first three coefficients are actually the same.

Let us introduce the series

$$\begin{aligned} y_2 = \tilde{X}(\tilde{Q}(x)^2) = & x^2 + 1488 x^3 + 2053632 x^4 + 2621653632 x^5 \\ & + 3244440682476 x^6 + \frac{19627900112688192}{5} x^7 + \frac{23401843163094440736}{5} x^8 \\ & + \frac{193179165341208747259392}{35} x^9 + \dots \end{aligned} \quad (295)$$

This series (295) is solution of the Schwarzian equation (174), with $W(x)$ given by (283), and is also solution of

$$2 \cdot F(y_2) = F(x) \cdot \frac{\partial y_2}{\partial x}, \quad (296)$$

i.e.

$$2 \cdot (y_2 - 744 y_2^2 - 393768 y_2^3) = (x - 744 x^2 - 393768 x^3) \cdot \frac{\partial y_2}{\partial x}, \quad (297)$$

and one also has:

$$\tilde{Q}(y_2(x)) = \tilde{Q}(x)^2. \quad (298)$$

Let us introduce the *one-parameter* series

$$\begin{aligned} y_2^{(a)} = y(a, y_2) = \tilde{X}(a \cdot \tilde{Q}(x)^2) = & a \cdot x^2 + 1488 a \cdot x^3 \\ & - 24 \cdot a \cdot (31 a - 85599) \cdot x^4 - 35712 \cdot a \cdot (62 a - 73473) \cdot x^5 \\ & + 36 \cdot a \cdot (9907 a^2 - 130673184 a + 90254015568) \cdot x^6 \\ & + \frac{160704}{5} \cdot a \cdot (49535 a^2 - 262999040 a + 122399922528) \cdot x^7 + \dots \end{aligned} \quad (299)$$

This one-parameter series (299) is solution of the Schwarzian equation (174), with $W(x)$ given by (283). It is also solution of

$$\tilde{Q}(y_2^{(a)}(x)) = a \cdot \tilde{Q}(x)^2, \quad (300)$$

and also solution of

$$2 \cdot F(y_2^{(a)}) = F(x) \cdot \frac{\partial y_2^{(a)}}{\partial x} = 2 \cdot a \cdot \frac{\partial y_2^{(a)}}{\partial a}. \quad (301)$$

where

$$F(x) = x - 744 x^2 - 393768 x^3 = x \cdot (1 - p \cdot x) \cdot (1 - q \cdot x), \quad (302)$$

with:

$$p = 372 + 6 \cdot 14782^{1/2}, \quad q = 372 - 6 \cdot 14782^{1/2}, \quad (303)$$

Let us denote

$$\alpha = \frac{1}{2} \cdot \frac{p+q}{q-p} = -\frac{31}{14782} \cdot 14782^{1/2} = -0.25497 \dots \quad (304)$$

Following the previous calculations in section (9.2), one easily finds that the “nome-like” series (286) reads:

$$\begin{aligned} \tilde{Q}(x) = & \frac{x \cdot (1 - p \cdot x)^{p/(q-p)}}{(1 - q \cdot x)^{q/(q-p)}} = \frac{x}{((1 - p \cdot x) \cdot (1 - q \cdot x))^{1/2}} \cdot \left(\frac{1 - p \cdot x}{1 - q \cdot x} \right)^\alpha \\ = & x + 744 x^2 + 750420 x^3 + 753621408 x^4 + 782312864472 x^5 \\ & + \frac{4097211834177216}{5} x^6 + \frac{4331866321367059104}{5} x^7 + \dots \end{aligned} \quad (305)$$

This “nome-like” series (305) is *actually* D -finite. It is solution of the order-one linear differential operator ($\theta = x \cdot D_x$ is the homogeneous derivative):

$$\begin{aligned}\mathcal{L}_1 &= F(x) \cdot D_x - 1 = (x - 744x^2 - 393768x^3) \cdot D_x - 1 \\ &= (1 - 744x - 393768x^2) \cdot \theta - 1.\end{aligned}\quad (306)$$

Do note, however, that this “nome-like” series (305) is *not* globally bounded. The radius of convergence of the “nome-like” series (305) is $1/p$, with p given by (303):

$$R = \frac{1}{p} = \frac{14782^{1/2}}{65628} - \frac{31}{32814} = 0.0009078632370 \dots \quad (307)$$

This “nome” series (305) is D -finite, with a finite radius of convergence, *but it is not globally bounded*. Note that $\tilde{X}(x)$ is only a *differentially algebraic function*.

Note that the order-one linear differential operator \mathcal{L}_1 , given by (306), is *not globally nilpotent* [83]. The corresponding p -curvatures are null (or nilpotent that is the same for order-one linear differential operators) for the following primes:

$$3, 11, 13, 17, 23, 31, 47, 61, 73, 79, 89, 101, \dots \quad (308)$$

but non-zero for the following primes:

$$5, 7, 19, 29, 37, 41, 43, 53, 59, 67, 71, \dots \quad (309)$$

Note that, since $14782 = 2 \cdot 19 \cdot 389$, we could have expected that one does not see the transcendence of the “nome” mod. 19, the “nome” reducing to an algebraic function (see (303), (304)), and thus one could expect a zero p -curvature. This is not the case.

Note that the exponent of the “nome-like” series (305), at the singularity $x = 1/p$, is

$$\frac{p}{q-p} = -\frac{1}{2} - \frac{31}{14782^{1/2}} = -0.7549735291 \dots \quad (310)$$

which is not a rational number. This rules out the fact that the order-one linear differential operator (306) could be globally nilpotent [83].

Let us consider the simplest example of series $y(a, x)$, namely the (involutive) series (284) for $a = -1$:

$$\begin{aligned}y(-1, x) &= -x - 1488x^2 - 2214144x^3 - 3099337344x^4 - 4030574598144x^5 \\ &\quad - \frac{23640158283604992}{5}x^6 - \frac{23310435220175683584}{5}x^7 \\ &\quad - \frac{20590422517553304526848}{7}x^8 + \frac{12494610391145690921435136}{7}x^9 + \dots\end{aligned}\quad (311)$$

Calculating the first fifty coefficients of this series, one can see that this (involutive) series is *not globally bounded*.

Remark 9.2: Following subsection (9.2.1), the generalization to two-parameter series can be performed on this last polynomial example, *mutatis mutandis*.

10. Comments and speculations on differentially algebraic series.

We have displayed miscellaneous series solutions of Schwarzian equations (and thus having a compositional property [30, 31]), which can be seen to be, or to generalize, *modular correspondences* [76]. We remark that we have the following situation:

we have series depending on *one* parameter (sometimes *two parameters* for slightly “academical” examples like in subsection (8.1)), which reduce to series with *integer* coefficients for an *infinite set* of values of the parameter(s), namely the *integer values*[‡]. These one-parameter series are *generically only differentially algebraic*, even for *integer* values of the parameter (where they are probably not even *D*-finite, see for instance (161)). In contrast, and remarkably, when the parameter is a *N*-th root of unity, the generically differentially algebraic one-parameter series become *algebraic functions*. We thus have an *infinite number* of *algebraic functions*.

It is interesting to note that a totally and utterly similar situation have been seen to occur in other very interesting situations in physics, or enumerative combinatorics. Along this line, differentially algebraic series with *integer* coefficients[¶] exist, and correspond to remarkable solutions of *differentially algebraic* equations in physics, or enumerative combinatorics, like *λ -extensions of Ising correlation functions* [84, 85], or solution of a differentially algebraic Tutte equation [55]. We have an *infinite set* of differentially algebraic series with *integer* coefficients that are *not D*-finite [55, 84, 85]. We also have the occurrence of an *infinite number* of *algebraic series* for an *infinite set* of Tutte-Beraha values of the λ parameter. Note that these selected values can also be seen as *N*-th root of unity situation.

At first sight, these Tutte-Beraha examples [55], or λ -extension of correlation functions of the Ising model [84, 85], *are not related to Schwarzian equations* with their composition function properties[†]. Is it possible that such differentially algebraic series could also reduce (in a more or less involved way ...) to exact decompositions like $X(\omega \cdot Q^n(x))$, that we found systematically through this paper, since many of the results of this paper are, in fact, consequences of such exact decompositions ?

- One motivation of this paper was to understand the very nature of the one-parameter series $y(a, x)$: we have seen that this series cannot be solution of an order-*N* linear differential operator (for some integer *N* independent of the parameter *a*) with coefficients polynomials in *x* and in the parameter *a*.

- The relation between the Schwarzian equations (such that $W(x) = -1/2/x^2 + \dots$, see [30, 31]), and *modular correspondences* was also an important motivation. The solutions of the Schwarzian equations are larger than just the (infinite) set of “modular correspondences”, *precisely* because of the occurrence of one-parameter series $y(a, x)$. Along this line we have first seen that the solution of the Schwarzian equations can actually correspond to series with *more than one parameter*. Modular correspondences are associated with modular curves and modular forms [14, 15, 16]. Consequently, another question was to know if one can generalize these concepts *beyond* the elliptic curves and modular forms framework.

We have also shown, with very simple (polynomial) examples for the function $F(x)$, that these structures can actually be generalized *far beyond* the elliptic curve (modular curve, Shimura curves, modular form, automorphic form) framework. Along this line, a first polynomial example (9.2) provides an example of *one-parameter series* $y(a, x)$, *algebraic for any value of the parameter*. We also found that the equivalent of the nome is a simple algebraic function (square root of a rational function). With that

[‡] More generally, for *rational values* of the parameters we have globally bounded differentially algebraic series.

[¶] Not simply reducible to ratio of globally bounded *D*-finite series, or composition of globally bounded *D*-finite series.

[†] These λ -extension of Ising correlation functions are solutions of *Painlevé equations* [84, 85].

example one also understands why it can be extremely hard to see that some series are algebraic, even if the algebraic function to guess is of a quite simple form. Furthermore, a “truncated” example (9.3) shows that the “modular equation-like” series (see for instance (295), (311)) can actually be *non globally bounded*. The “nome-like” series is a *non globally bounded*, but *still D-finite*, series (see (306)), the corresponding linear differential operator being *non globally nilpotent*.

11. Conclusion

This paper provides a simple, and pedagogical, illustration of exact non-linear symmetries in physics (exact representations of the renormalization group transformations like the *Landen transformation* for the square Ising model [32, 33], ...) and is a strong incentive to discover more *differentially algebraic* equations involving fundamental symmetries, and to develop more differentially algebraic series analysis in physics [55, 56], beyond examples like the full susceptibility of the square-lattice Ising model [56, 84, 85, 92, 93].

In this paper we first focused, essentially, on identities relating the *same* hypergeometric function with *two different* algebraic pullback transformations related by modular equations. This corresponds to the “classical” *modular forms* [19] (resp. automorphic forms) that emerged so many times in physics [41, 42, 43]: these algebraic transformations can be seen as simple illustrations of exact representations of the renormalization group of some Yang-Baxter integrable models [32, 33, 73]. These transformations are seen to be solutions of some Schwarzian relation.

The Schwarzian relation is seen to “encapsulate”, in one differentially algebraic (Schwarzian) equation, all the *modular forms* and *modular equations* of the theory of elliptic curves. The Schwarzian condition can thus be seen as some quite fascinating “pandora box”, which encapsulates an *infinite number* of highly remarkable modular equations, and a whole “universe” of *Belyi-maps*‡. It is however important to underline that these Schwarzian conditions are actually richer than just elliptic curves, and go beyond†† “simple” restrictions [91] to pullbacked ${}_2F_1$ hypergeometric functions. In a more general perspective, such Schwarzian conditions occur in Malgrange’s pseudo-group approach [65, 66, 67, 72] of \mathcal{D} -enveloppes. At this level of mathematical abstraction, the question of a *modular correspondence interpretation* of these “Schwarzian” series was clearly an open question. This paper sheds some light on this open question. It shed some light on the very nature of the one-parameter series solution of the Schwarzian equation, which is *not* generically a modular correspondence series, but *actually reduces* to an *infinite set of modular correspondence series* for an infinite set of (N -th root of unity) values of the parameter. This paper also provides (polynomial) examples that are very similar to modular correspondence series, but are far beyond the elliptic curves framework.

Acknowledgments: We would like to thank A. Bostan and G. Casale, for very fruitful discussions. We thank A. Bostan for some p -curvature calculations. This work has been performed without any ERC, ANR, PES or MAE financial support.

‡ Belyi-maps [86, 87, 88, 89, 90] are central to Grothendieck’s program of “dessins d’enfants”.

†† See the two Heun functions given by (164) in [77].

Appendix A. Miscellaneous modular correspondences.

• Let us consider

$$\begin{aligned} \tilde{X}(\tilde{Q}(x)^7) = & x^7 + 5208x^8 + 16877196x^9 + 43972589024x^{10} \\ & + 101156052918270x^{11} + 215029151897268240x^{12} + \dots \end{aligned} \quad (\text{A.1})$$

which is solution of a modular equation $\Gamma_7(x, y) = \Gamma_7(y, x) = 0$ that we will not write here, but can easily be obtained from its rational parametrization [19]:

$$x = \frac{t}{(t^2 + 13t + 49) \cdot (t^2 + 5t + 1)^3}, \quad y = \frac{t^7}{(t^2 + 13t + 49) \cdot (t^2 + 245t + 2401)^3}.$$

This series (A.1) is solution of an order-eight linear differential operator $L_8 = L_1 \oplus L_7$, which is the direct sum of an order-one linear differential operator L_1 with a rational function solution of the form

$$x \cdot \frac{p_6(x)}{(11356800389480448000000x^2 + 34848505552896000x + 1)^3}, \quad (\text{A.2})$$

where $p_6(x)$ is a polynomial of degree six, and an order-seven irreducible linear differential operator L_7 .

• Let us also consider

$$\tilde{X}(\tilde{Q}(x)^8) = x^8 + 5952x^9 + 21502368x^{10} + 61310179840x^{11} + \dots \quad (\text{A.3})$$

which is solution of a modular equation $\Gamma_8(x, y) = \Gamma_8(y, x) = 0$ that we will not write here, but can easily be obtained from its rational parametrization [19]:

$$\begin{aligned} x &= \frac{t \cdot (t+4)^2 \cdot (t+8)}{(t^4 + 16t^3 + 80t^2 + 128t + 16)^3}, \\ y &= \frac{t^8 \cdot (t+4) \cdot (t+8)^2}{(t^4 + 256t^3 + 5120t^2 + 32768t + 65536)^3}. \end{aligned} \quad (\text{A.4})$$

The polynomial, associated with the modular equation $\Gamma_8(x, y) = 0$, is of degree 12 in y (resp. in x). This series (A.3) is solution of an order-twelve linear differential operator $L_{12} = L_1 \oplus L_{11}$, which is the direct sum of an order-one linear differential operator L_1 with a rational function solution of the form

$$x \cdot \frac{p_{11}(x)}{p_4(x)^3}, \quad (\text{A.5})$$

where $p_{11}(x)$ is a polynomial of degree eleven, and where $p_4(x)$ reads

$$\begin{aligned} & 1080060886113159937649308593750000x^4 - 826335556188178615474500000000x^3 \\ & - 15705521635909735050750000x^2 + 8041801037378436000x - 1, \end{aligned} \quad (\text{A.6})$$

and an order-eleven linear differential operator L_{11} . The other *algebraic* solution series of $\Gamma_8(x, y) = 0$ are the compositional inverse of series (A.3), namely

$$\begin{aligned} & \omega \cdot x^{1/8} - 744 \cdot \omega^2 \cdot x^{2/8} + 356652\omega^3 \cdot x^{3/8} - 140361152 \cdot \omega^4 \cdot x^{4/8} \\ & + 49336682190 \cdot \omega^5 \cdot x^{5/8} - 16114625669088 \cdot \omega^6 \cdot x^{6/8} + \dots \end{aligned} \quad (\text{A.7})$$

where $\omega^8 = 1$, together with

$$\begin{aligned} & -x^2 - 1488x^3 - 2055120x^4 - 2864378368x^5 - 4071821465856x^6 \\ & - 5900566305239040x^7 - 8695398352685449216x^8 + \dots \end{aligned} \quad (\text{A.8})$$

$$\begin{aligned} \omega \cdot x^{1/2} &- 744 \cdot \omega^2 \cdot x^{2/2} + 356280 \cdot \omega^3 \cdot x^{3/2} - 139807616 \cdot \omega^4 \cdot x^{4/2} \\ &+ 48938964576 \cdot \omega^5 \cdot x^{5/2} + \dots \end{aligned} \quad (\text{A.9})$$

- The (algebraic) series

$$\tilde{X}(\tilde{Q}(x)^{11}) = x^{11} + 8184x^{12} + 38699100x^{13} + 138966918112x^{14} + \dots \quad (\text{A.10})$$

$$x \cdot \frac{p_{11}}{p_1^3}, \quad (\text{A.11})$$
$$\begin{aligned}
& 1577314437358442913340940353536000000000000 x^4 \\
& - 496864268553728774541064273920000000000 x^3 \\
& + 45688143672322270430861721600000000 x^2 \\
& + 98823634118413525094400000 x + 1, \tag{A.12}
\end{aligned}$$

- The (algebraic) series

$$\tilde{X}\left(\tilde{Q}(x)^{12}\right)=x^{12}+8928 x^{13}+45538416 x^{14}+174773255424 x^{15}+\cdots \quad(\text { A.13 })$$

$$x = t \cdot \frac{(t+2)^3(t+3)^4(t+4)^3(t+6)}{(t^2+6t+6)^3 \cdot p_6^3}, \quad y = t^{12} \cdot \frac{(t+2)(t+3)^3(t+4)^4(t+6)^3}{(t^2+12t+24)^3 \cdot q_6^3},$$
$$\begin{aligned} p_6 &= t^6 + 18t^5 + 126t^4 + 432t^3 + 732t^2 + 504t + 24, \\ q_6 &= t^6 + 252t^5 + 4392t^4 + 31104t^3 + 108864t^2 + 186624t + 124416. \end{aligned} \quad (\text{A.14})$$
$$x \cdot \frac{p_{23}}{(6549518250000 x^2 - 2835810000 x + 1)^3 \cdot p_6^3} \quad (\text{A.15})$$

[†] See Appendix I in [44], which is the unabridged arXiv version of [43].

where p_{23} is a polynomial of degree 23, and where p_6 is the polynomial

$$\begin{aligned} & 42889619864187195342544128412237640625000000000000 x^6 \\ & + 3869372376492639837782614434923625000000000000 x^5 \\ & + 34904627315764077727184412247908187500000000 x^4 \\ & + 1007059405271040783775694468925000000000 x^3 \\ & + 280179539493990596285512318134750000 x^2 \\ & - 22804995243537595825782822000 x + 1, \end{aligned} \quad (\text{A.16})$$

and an order-23 linear differential operator L_{23} . The other (algebraic) series are, respectively, the compositional inverse of series (A.13), namely

$$\begin{aligned} & \omega \cdot x^{1/12} - 744 \cdot \omega^2 \cdot x^{2/12} + 356652 \cdot \omega^3 \cdot x^{3/12} - 140361152 \cdot \omega^4 \cdot x^{4/12} \\ & + 49336682190 \cdot \omega^5 \cdot x^{5/12} - 16114625669088 \cdot \omega^6 \cdot x^{6/12} \\ & + 4999042477430456 \cdot \omega^7 \cdot x^{7/12} + \dots \end{aligned} \quad (\text{A.17})$$

where $\omega^{12} = 1$, together with

$$-x^3 - 2232x^4 - 3911868x^5 - 6380015304x^6 - 10139549171670x^7 + \dots \quad (\text{A.18})$$

which is nothing but $y_3(y_1(x)) = y_1(y_3(x))$ (with y_3 given by (55), and y_1 given by (81)), and

$$\begin{aligned} & \omega \cdot x^{1/3} - 744 \cdot \omega^2 \cdot x^{2/3} - 356652 \cdot x + 140361400 \cdot \omega \cdot x^{4/3} \\ & - 49337051214 \cdot \omega^2 \cdot x^{5/3} - 16114891018176 \cdot x^{6/3} \\ & + 4999181715881876 \cdot \omega \cdot x^{7/3} + \dots \end{aligned} \quad (\text{A.19})$$

where $\omega^3 = 1$,

$$\omega \cdot x^{4/3} + 992 \cdot \omega \cdot x^{7/3} - 744 \cdot \omega^2 \cdot x^{8/3} + 1123568 \cdot \omega \cdot x^{10/3} + \dots \quad (\text{A.20})$$

where $\omega^3 = -1$, and

$$\omega \cdot x^{3/4} - 744 \cdot \omega^2 \cdot x^{6/4} + 558 \cdot \omega \cdot x^{7/4} + 356652 \cdot \omega^3 \cdot x^{9/4} + \dots \quad (\text{A.21})$$

where $\omega^4 = 1$. This gives $1 + 1 + 12 + 3 + 3 + 4 = 24$ algebraic series, solutions of $\Gamma_{12} = 0$ and L_{24} .

Remark 3.6: Recalling the algebraic series y_3 , given by (55), and the algebraic series y_1 given by (81), one can see that the algebraic series (A.18) is *nothing but*:

$$y_3(y_1(x)) = y_1(y_3(x)) = -x^3 - 2232x^4 - 3911868x^5 + \dots \quad (\text{A.22})$$

- The (algebraic) series

$$\begin{aligned} \tilde{X}(\tilde{Q}(x)^{16}) &= x^{16} + 11904x^{17} + 78431040x^{18} + 378584548352x^{19} \\ &+ 1496557573544352x^{20} + \dots \end{aligned} \quad (\text{A.23})$$

is solution of a *modular equation* $\Gamma_{16}(x, y) = \Gamma_{16}(y, x) = 0$, that we will not write here, but can easily be obtained from its rational parametrization [19]:

$$x = t \cdot \frac{(t+2)^4(t+4)(t^2+4t+8)}{p_8^3}, \quad y = t^{16} \cdot \frac{(t+2)(t+4)^4(t^2+4t+8)}{q_8^3},$$

where:

$$\begin{aligned} p_8 &= t^8 + 16t^7 + 112t^6 + 448t^5 + 1104t^4 + 1664t^3 + 1408t^2 + 512t + 16, \\ q_8 &= t^8 + 256t^7 + 5632t^6 + 53248t^5 + 282624t^4 + 917504t^3 + 1835008t^2 \\ &+ 2097152t + 1048576. \end{aligned} \quad (\text{A.24})$$

The polynomial in the modular equation $\Gamma_{16}(x, y) = 0$ is of degree 24 in y (resp. in x), and, thus, has twenty four algebraic solution series, corresponding to the series (A.23), the compositional inverse of series (A.23), namely

$$\begin{aligned} \omega \cdot x^{1/16} &- 744 \cdot \omega^2 \cdot x^{2/16} + 356652 \cdot \omega^3 \cdot x^{3/16} \\ &- 140361152 \cdot \omega^4 \cdot x^{4/16} + \dots \end{aligned} \quad (\text{A.25})$$

where $\omega^{16} = 1$, together with

$$\begin{aligned} -x^4 &- 2976x^5 - 6322896x^6 - 11838151424x^7 - 20872495229904x^8 \\ &- 35647177059836928x^9 + \dots \end{aligned} \quad (\text{A.26})$$

and

$$\begin{aligned} \omega \cdot x^{1/4} &- 744 \cdot \omega^2 \cdot x^{2/4} + 356652 \cdot \omega^3 \cdot x^{3/4} \\ &- 140361152 \cdot \omega^4 \cdot x^{4/4} + \dots \end{aligned} \quad (\text{A.27})$$

where $\omega^4 = -1$, and:

$$\omega \cdot x - 744 \cdot \omega \cdot (\omega - 1) \cdot x^2 + \dots \quad (\text{A.28})$$

where $\omega^2 = -1$. The (algebraic) series solutions of the modular equation $\Gamma_{16}(x, y) = 0$ are solution of an order-24 linear differential operator $L_{24} = L_1 \oplus L_{23}$, which is the direct sum of an order-23 linear differential operator L_{23} , and an order-one linear differential operator L_1 , with a rational function solution

$$x \cdot \frac{p_{23}}{p_8^3}, \quad (\text{A.29})$$

where p_{23} is a polynomial of degree 23, and where p_8 is the polynomial

$$\begin{aligned} &15926143836920796849094002857387135460968690480161221686575776100158691406250000x^8 \\ &- 6042818923606714182438083804301870179528875596947614517314453125000000000000x^7 \\ &+ 49006987053737646413653547579882802471367855785728981543291015625000000000x^6 \\ &+ 467218903177861854107002271749529441241375461552376767332031250000000000x^5 \\ &+ 81580198367732340212612911642019252294658707587093110574218750000x^4 \\ &+ 7361546087090590150064981160492829297036925882551350000000000x^3 \\ &+ 259399171372225204966661002550162965440584749500000x^2 \\ &- 64670563924749466394147714711210760000x + 1. \end{aligned} \quad (\text{A.30})$$

Remark 3.7: Recalling the algebraic series y_4 , given by (79), and the algebraic series y_1 given by (81), one can see that the algebraic series (A.26) is *nothing† but*:

$$y_1(y_4(x)) = -x^4 - 2976x^5 - 6322896x^6 + \dots \quad (\text{A.31})$$

We thus have $1 + 1 + 16 + 4 + 2 = 24$ algebraic solutions of the modular equation $\Gamma_{16}(x, y) = 0$, and also solutions of L_{24} .

- The (algebraic) series

$$\begin{aligned} \tilde{X}(\tilde{Q}(x)^{18}) &= x^{18} + 13392x^{19} + 98198568x^{20} + 522607392000x^{21} \\ &+ 2259156547520244x^{22} + \dots \end{aligned} \quad (\text{A.32})$$

† In contrast $y_4(x) = y_4(y_1(x))$.

$$\begin{aligned} x &= t \cdot \frac{(t+2)^9 \cdot (t+3)^2 \cdot (t^2+3t+3)^2 (t^2+6t+12)}{(t^3+6t^2+12t+6)^3 \cdot p_9^3}, \\ y &= t^{18} \cdot \frac{(t+2)^2 \cdot (t+3)^9 \cdot (t^2+3t+3) (t^2+6t+12)^2}{(t^3+12t^2+36t+36)^3 \cdot q_9^3}, \end{aligned} \quad (\text{A.33})$$
$$\begin{aligned}
p_9 &= t^9 + 18t^8 + 144t^7 + 666t^6 + 1944t^5 + 3672t^4 \\
&\quad + 4404t^3 + 3096t^2 + 1008t + 24, \\
q_9 &= t^9 + 252t^8 + 4644t^7 + 39636t^6 + 198288t^5 + 629856t^4 \\
&\quad + 1294704t^3 + 1679616t^2 + 1259712t + 419904.
\end{aligned} \tag{A.34}$$
$$\begin{aligned} & \omega \cdot x^{1/18} - 744 \cdot \omega^2 \cdot x^{2/18} + 356652 \cdot \omega^3 \cdot x^{3/18} \\ & - 140361152 \cdot \omega^4 \cdot x^{4/18} + \dots \end{aligned} \quad (\text{A.35})$$
$$\begin{aligned} \omega \cdot x^2 &+ 1488 \cdot \omega \cdot x^3 + (744 + 2055120 \cdot \omega) \cdot x^4 \\ &+ (2214144 + 2864378368 \cdot \omega) \cdot x^5 + \dots \end{aligned} \quad (\text{A.36})$$
$$\begin{aligned} \omega \cdot x^{1/2} &- 744 \cdot \omega^2 \cdot x^{2/2} + (356652 \cdot \omega^3 + 372 \cdot \omega) \cdot x^{3/2} \\ &+ (139807616 \omega^2 + 140361152) \cdot x^{4/2} + \dots \end{aligned} \quad (\text{A.37})$$
$$\omega \cdot x^{2/9} - 744 \cdot \omega^2 \cdot x^{4/9} + 356652 \cdot \omega^3 \cdot x^{6/9} + \dots \quad (\text{A.38})$$
$$\omega \cdot x^{9/2} + 3348 \cdot \omega \cdot x^{11/2} + 7735986 \cdot \omega \cdot x^{13/2} + \dots \quad (\text{A.39})$$
$$x \cdot \frac{p_{35}}{p_3^3 \cdot p_9^3}, \quad (\text{A.40})$$
$$1879994705688000000000 x^3 - 224179462188000000 x^2 + 151013228706000 x - 1,$$
$$\begin{aligned} & 141600617083186841426749541059379178266125496444877735060776646144000000000000000000000 x^9 \\ & - 91940358193098820927255075706021981712433442298247865135275206912000000000000000000000 x^8 \\ & + 2357512764312464299933742109740167360806718661721497394039023769600000000000000000000 x^7 \\ & + 47239695875311014088873100371849646543690662098121255705065139200000000000000000000 x^6 \end{aligned}$$

$$\begin{aligned}
& + 97654415151135929389144490831366836006678881009953401828037760000000000000000 x^5 \\
& + 140388149200703815304879817897188959087877294047069901286720000000000000 x^4 \\
& + 6390980152781882840426709358572754975540778747694537696000000000 x^3 \\
& + 7559858588621896366536922746878187128255472000000 x^2 \\
& + 3443855962300764146093216928806375326182000 x - 1.
\end{aligned} \tag{A.41}$$

Appendix B. Beyond pullbacked ${}_2F_1$ hypergeometric functions: a selected Heun function.

Let us show that the results displayed on the classical modular curves and their associated modular forms with pullbacked ${}_2F_1$ hypergeometric functions, also work on Shimura curves [75] and their associated automorphic forms [10] with a Heun function *which cannot be reduced to pullbacked ${}_2F_1$ hypergeometric functions*.

Recalling[†] Krammer's counterexample to Dwork's conjecture [78, 79, 80], let us consider the Heun function $Heun(81, 1/2, 1/6, 1/3, 1/2, 1/2; x)$ which is solution of an order-two linear differential operator L_2 which is *globally nilpotent* [83]. The series expansion of this Heun function is *not globally bounded* [43, 44]. Let us introduce the following function $F(x)$:

$$F(x) = x^{1/2} \cdot \left(1 - \frac{x}{81}\right)^{1/2} \cdot (1-x)^{1/2} \cdot HeunG\left(81, \frac{1}{2}, \frac{1}{6}, \frac{1}{3}, \frac{1}{2}, \frac{1}{2}; x\right)^2, \tag{B.1}$$

or, more simply, the following (non globally bounded) series:

$$\begin{aligned}
81 \cdot F(x)^2 &= x \cdot (81-x) \cdot (1-x) \cdot HeunG\left(81, \frac{1}{2}, \frac{1}{6}, \frac{1}{3}, \frac{1}{2}, \frac{1}{2}; x\right)^4 \\
&= 81x - 78x^2 - \frac{137}{81}x^3 - \frac{3892}{6561}x^4 - \frac{44495}{177147}x^5 - \frac{1900594}{14348907}x^6 + \dots
\end{aligned} \tag{B.2}$$

Let us consider the Schwarzian equation associated with the order-two linear differential operator L_2 . The corresponding function $W(x)$ reads (see subsection (1.2)):

$$W(x) = -\frac{35x^4 - 3680x^3 + 244242x^2 - 244944x + 177147}{72 \cdot x^2 \cdot (x-1)^2 \cdot (x-81)^2}. \tag{B.3}$$

One can actually verify that $W(x)$, given by the *rational function* (B.3), can actually also be written *in terms of the Heun function* (B.1):

$$W(x) = \frac{F''(x)}{F(x)} - \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)}\right)^2 = \left(\frac{F'(x)}{F(x)}\right)' + \frac{1}{2} \cdot \left(\frac{F'(x)}{F(x)}\right)^2. \tag{B.4}$$

Note that introducing a “nome”

$$\frac{Q(x)'}{Q(x)} = \frac{1}{F(x)} \quad \text{or:} \quad Q(x) = \exp\left(\int^x \frac{dx}{F(x)}\right), \tag{B.5}$$

relation (B.4) is nothing but relation (25), namely:

$$W(x) = -\{Q(x), x\} - \frac{1}{2 \cdot Q(x)^2} \cdot \left(\frac{dQ(x)}{dx}\right)^2. \tag{B.6}$$

A one-parameter series $y(a, x)$ is actually solution[†] of the Schwarzian equation

$$W(x) - W(y(x)) \cdot y'(x)^2 + \{y(x), x\} = 0, \tag{B.7}$$

[†] See also subsection 2.3 of [83].

[†] Note that there are no solutions of the form $a \cdot x^n + \dots$ with $n \geq 2$, since $W(x)$ is not of the form $-1/2/x^2 + \dots$. The series expansion of $W(x)$ reads $-3/8/x^2 + \dots$

with $W(x)$ given by (B.3):

$$y(a, x) = a \cdot x - \frac{26}{81} \cdot a \cdot (a-1) \cdot x^2 + \frac{1}{6561} \cdot a \cdot (a-1) \cdot (243a - 1109) \cdot x^3 \\ - \frac{2}{3720087} \cdot a \cdot (a-1) \cdot (4013a^2 - 62326a + 201028) \cdot x^4 + \dots \quad (\text{B.8})$$

One can easily verify the functional equation:

$$a \cdot F\left(y(a, x)\right)^2 = F(x)^2 \cdot \frac{dy(a, x)}{dx}, \quad (\text{B.9})$$

or:

$$a^{1/2} \cdot F\left(y(a, x)\right) = F(x) \cdot \frac{dy(a, x)}{dx}. \quad (\text{B.10})$$

One can also verify the following compositional formula:

$$y\left(a, y(a', x)\right) = y(a a', x). \quad (\text{B.11})$$

Let us introduce the two $a \rightarrow 0$ and $a \rightarrow \infty$ limits of the one-parameter series (B.8):

$$\tilde{Q}(x) = \lim_{a \rightarrow 0} \frac{y(a, x)}{a} = x + \frac{26}{81} x^2 + \frac{1109}{86561} x^3 + \frac{402056}{3720087} x^4 \\ + \frac{2565526}{33480783} x^5 + \frac{471402140}{8135830269} x^6 + \dots \quad (\text{B.12})$$

and:

$$\tilde{X}(x) = \lim_{a \rightarrow \infty} y\left(a, \frac{x}{a}\right) = x - \frac{26}{81} x^2 + \frac{1}{27} x^3 - \frac{8026}{3720087} x^4 \\ + \frac{38603}{301327047} x^5 - \frac{3200}{301327047} x^6 + \dots \quad (\text{B.13})$$

One verifies that the one-parameter series (B.8) is *actually of the form*:

$$y(a, x) = \tilde{X}\left(a \cdot \tilde{Q}(x)\right). \quad (\text{B.14})$$

From the exact decomposition (B.14), together with the fact that the one-parameter series (B.8) is such that $y(1, x) = 1$, one deduces immediately that the series (B.13) is actually the *compositional inverse* of the series (B.12).

Appendix C. Very simple polynomial examples for $F(x)$.

Let us display some very simple examples for $F(x)$, and the corresponding one-parameter functions $y(a, x)$, solutions of the Schwarzian equation (17).

• For $F(x) = x \cdot (1 + px)$, one has $W(x) = -1/2/x^2/(1 + px)^2$, and a one-parameter function $y(a, x)$, solution of the Schwarzian equation (17) with that $W(x)$ reads:

$$y(a, x) = \frac{a \cdot x}{(1 + px) - a \cdot p \cdot x}. \quad (\text{C.1})$$

It is straightforward to see that $y(a, x)$ can be written $y(a, x) = \tilde{X}(a \cdot \tilde{Q}(x))$, where:

$$\tilde{X}(x) = \frac{x}{1 - p \cdot x} \quad \text{and:} \quad \tilde{Q}(x) = \frac{x}{1 + p \cdot x}. \quad (\text{C.2})$$

• For $F(x) = x + p$, one has $W(x) = -1/(x + p)^2/2$, and a one-parameter function $y(a, x)$, solution of $F(y(a, x)) = F(x) \cdot \frac{\partial(a, x)}{\partial x}$ and of the Schwarzian equation (17) with that $W(x)$ reads (p is fixed):

$$y(a, x) = a \cdot (x + p) - p. \quad (\text{C.3})$$

• For $F(x) = x^2$, one has[†] $W(x) = 0$, and a one-parameter function $y(a, x)$, solution of $F(y(a, x)) = F(x) \cdot \frac{\partial(a, x)}{\partial x}$ and of the Schwarzian equation (17) with $W(x) = 0$ reads:

$$y(a, x) = \frac{x}{1 - \ln(a) \cdot x}. \quad (\text{C.4})$$

• For $F(x) = p$, one has $W(x) = 0$, the one-parameter function $y(a, x)$, solution of $F(y(a, x)) = F(x) \cdot \frac{\partial(a, x)}{\partial x}$ and of the Schwarzian equation (17) with $W(x) = 0$ reads (p is fixed):

$$y(a, x) = x + p \cdot \ln(a). \quad (\text{C.5})$$

• For $F(x) = p \cdot x$, one has[¶] $W(x) = -1/2/x^2$, and the one-parameter function $y(a, x)$, solution of $F(y(a, x)) = F(x) \cdot \frac{\partial(a, x)}{\partial x}$ and of the Schwarzian equation (17) with that $W(x)$ reads (p is fixed):

$$y(a, x) = a^p \cdot x. \quad (\text{C.6})$$

All these one-parameter functions (C.1), (C.3), (C.4), (C.5), (C.6) verify the composition rule:

$$y(a, y(a', x)) = y(a a', x). \quad (\text{C.7})$$

All these one-parameter functions (C.1), (C.3), (C.4), (C.5), (C.6) verify:

$$F(y(a, x)) = a \cdot \frac{\partial y(a, x)}{\partial a} = F(x) \cdot \frac{\partial y(a, x)}{\partial x}. \quad (\text{C.8})$$

- [1] S. Boukraa, S. Hassani and J-M.Maillard, *Noetherian mappings*, Physica **D 185**, (2003) pp. 3-44
- [2] S. Boukraa and J-M.Maillard, *Factorization properties of birational mappings*, Physica **A 220** (1995) 403-470.
- [3] M.P. Bellon, J-M. Maillard and C. Viallet, *Infinite discrete symmetry group for the Yang-Baxter equations, Vertex models*, Physics Letters **B 260**, (1991) pp. 87-100.
- [4] M.P. Bellon, J-M. Maillard and C-M. Viallet, *Quasi-Integrability of the sixteen vertex model*, Phys.Lett. **B 281**, (1992), pp. 315-319
- [5] P. Fatou, Sur les équations fonctionnelles, Journal of Symbolic Comutation **94**, (2019), pp.90-104.
- [6] J. F. Ritt, On the iteration of rational function, Trans. Amer. Math. Soc. **21** (1920), pp. 348-356.
- [7] P. Fatou, Sur les fonctions qui admettent plusieurs théorèmes de multiplication, C. R. acad. Sci. Paris, Sér. I Math **173**, (1921), pp. 571-573.

[†] More generally one has $W(x) = 0$ for $F(x) = q \cdot (x + p)^2$. For $W(x) = 0$ the solution of the Schwarzian equation(17) is a three-parameter solution: $y(x) = \frac{a x + b}{c x + d}$.

[¶] More generally when $W(x) = -1/2/x^2$ the solution of the Schwarzian equation (17) with that $W(x)$, is a three-parameter solution: $\ln(y(x)) = \frac{a \ln(x) + b}{c \ln(x) + d}$.

- [8] A. E. Eremenko, On some functional equations connected with iteration of rational functions, Leningrad Math. J. Vol. 1, (1990), pp. 905-919.
- [9] B. M. McCoy and J-M. Maillard, *The anisotropic Ising correlations as elliptic integrals: duality and differential equations*, J. Phys. **A 49**, Number 43, (2016) 434004 (24pp), Special Issue in honour of A. J. Guttmann, arXiv:1606.08796v4 [math-ph]
- [10] L. Ford, *Automorphic Functions*, (1929) AMS Chelsea Publishing.
- [11] P. F. Stiller, *Classical Automorphic Forms and Hypergeometric Functions*, Journ. of Number Theory, **28**, no. 2, 219-232, (1988).
- [12] J. McKay and A. Sebbar, *Fuchsian groups, automorphic functions and Schwarzians*, Math. Ann. **318**, (2000) pp.255-275.
- [13] J. McKay and A. Sebbar, *Fuchsian groups, Schwarzians, and theta functions*, C. R. Acad. Sci. Paris, **327**, Série I, (1998) pp.343-348.
- [14] N. Koblitz, *Introduction to Elliptic Curves and Modular Forms*, Graduate Texts in Mathematics, Second Edition, 1993, Springer
- [15] J. H. Bruinier, G. van der geer, G. Harder and D. Zagier, *The 1-2-3 of Modular Forms*, Universitext, 2008, Springer
- [16] G.E. Andrews, B.C. Berndt, *Ramanujan's Lost Notebook*, 2012, Part I, Part III, Part V, Springer
- [17] Henri Cohen and Fredrik Stromberg, *Modular Forms: A Classical Approach*, Graduate Studies in Mathematics, **179**, American Mathematical Society
- [18] Paul B. Garrett, *Holomorphic Hilbert Modular Forms*, Wadsworth and Brooks/Cole Advanced Books and Software, Pacific Grove, CA, 1990
- [19] R. Maier, *On rationally parametrized modular equations*, J. Ramanujan Math. Soc (2009), pp. 1-73, (2006), arXiv:0611041v3 [math.NT]
- [20] C. McMullen, *Amenability, Poincaré series and quasiconformal maps*, Invent. Math. **97** (1989), 95-127.
- [21] R. Fricke, F. Klein, *Vorlesungen über die Theorie der automorphen Funktionen*, 1-2, Teubner (1926)
- [22] J. Kollár, *Shafarevich maps and automorphic forms*, (1995), M. B. Porter Lectures, Princeton University Press
- [23] D.P. Dalzell, *Theory of the theta-Fuchsian functions*, London Math. Soc. pp. 539-558
- [24] Farkas, Hershel M. (2008). *Theta functions in complex analysis and number theory*, In Alladi, Krishnaswami (ed.). Surveys in Number Theory. Developments in Mathematics. **17**, Springer-Verlag. pp. 57-87.
- [25] B. Schoeneberg, (1974). *IX. Theta series". Elliptic modular functions*, Die Grundlehren der mathematischen Wissenschaften, **203**, Springer-Verlag. pp. 203-226.
- [26] <http://www.physics.fsu.edu/courses/Spring05/phy6938-02/decimation.pdf>
- [27] M.E. Fisher, *Renormalization group theory: Its basis and formulation in statistical physics*, Reviews of Modern Physics, **70**, No. 2, (1998) pp. 653-681
- [28] A. Bostan, S. Boukraa, S. Hassani, J-M. Maillard, J-A. Weil, N. Zenine and N. Abarenkova, *Renormalization, isogenies and rational symmetries of differential equations*, Advances in Mathematical Physics, Hindawi Pub. Volume 2010, ID 941560, 44 pages
- [29] Antonio Jiménez-Pastor, Veronika Pillwein, *A computable extension for D-finite functions: DD-finite functions*, Journal of Symbolic Computation **94**, 2019, pp. 90-104
- [30] Y. Abdelaziz and J-M. Maillard, *Modular forms, Schwarzian conditions, and symmetries of differential equations in physics*, (2017) J.Phys. **A 50**: Math. Theor. 215203
- [31] Y. Abdelaziz and J-M. Maillard, *Schwarzian conditions for linear differential operators with selected differential Galois groups*, (2017) J.Phys. **A 50**: Math. Theor. 465201 (33p).
- [32] S. Boukraa, S. Hassani, J-M. Maillard and N. Zenine, *Singularities of n-fold integrals of the Ising class and the theory of elliptic curves*, J. Phys. **A 40**: Math. Theor (2007) 11713-11748 <http://arxiv.org/pdf/math-ph/0706.3367>
- [33] S. Boukraa, S. Hassani, J.-M. Maillard and N. Zenine, *Landau singularities and singularities of holonomic integrals of the Ising class*, J. Phys. A: Math. Theor. **40** (2007) 2583-2614 and arXiv:math-ph/0701016v2
- [34] B. C. Berndt and H. H. Chan, *Ramanujan and the Modular j-Invariant*, Canad. Math. Bull. **42** (1999) 427-440
- [35] G.E. Andrews and B.C. Berndt, Chapter 17 pp. 373-393, in *Ramanujan's Lost Notebook*, Part I, 52005) Springer
- [36] H. H. Chan and M.-L. Lang, *Ramanujan's modular equations and Atkin-Lehner involutions*, Israel Journal of Mathematics, **103**, (1998) pp. 1-16.
- [37] C. Hermite, *Sur la théorie des équations modulaires*, Comptes Rendus Acad. Sci. Paris **49**, 16-24, 110-118, and 141-144, 1859 Oeuvres complètes, Tome II. Paris: Hermann, p. 61, 1912.

- [38] M. Hanna, *The Modular Equations*, Proc. London Math. Soc. **28**, 46-52, 1928.
- [39] F. Morain, *Calcul du nombre de points sur une courbe elliptique dans un corps fini: aspects algorithmiques*, Journal de Théorie des Nombres de Bordeaux, tome 7, (1995) pp.255-282 and <https://eudml.org/doc/247643>
- [40] Weisstein, Eric W. "Modular Equation." From MathWorld - A Wolfram Web Resource. <http://mathworld.wolfram.com/ModularEquation.html>
- [41] A. Bostan, S. Boukraa, S. Hassani, M. van Hoeij, J-M. Maillard, J-A. Weil, N. J. Zenine, *The Ising model: from elliptic curves to modular forms and Calabi-Yau equations*, J. Phys. **A 44**: Math. Theor. (2011) (43 pp) 045204 IOP Select, and arXiv: 1007.69804 v1 [math-ph] and hal-00684883, version 1
- [42] M. Assis, S. Boukraa, S. Hassani, M. van Hoeij, J-M. Maillard, B.M. McCoy *Diagonal Ising susceptibility: elliptic integrals, modular forms and Calabi-Yau equations*, J. Phys. **A 45**: Math. Theor. (2012) 075205, [32 pages]. IOP Select paper and Highlights of 2012 arXiv:arXiv:1110.1705v2 [math-ph]
- [43] A. Bostan, S. Boukraa, G. Christol, S. Hassani, J-M. Maillard, *Ising n-fold integrals as diagonal of rational functions and integrality of series expansions: integrality versus modularity*, (2012) J. Phys. A: Math. Theor. **46** 185202
- [44] A. Bostan, S. Boukraa, G. Christol, S. Hassani, J-M. Maillard, *Ising n-fold integrals as diagonal of rational functions and integrality of series expansions: integrality versus modularity (unabridged version)*, (2012) arXiv:1211.6031v1 [math-ph]
- [45] A.M. Legendre, *Traité des fonctions elliptiques*, vol. **3**, Paris, 1825-1828.
- [46] C.G.J. Jacobi, *Fundamenta Nova Theoriae Functionum Ellipticarum* Königsberg, 1829.
- [47] V. Ovsienko and S. Tabachnikov, *What is ... the schwarzian Derivative ?*, (2009), Notice of the AMS, **56**, pp. 34-36
- [48] A. Jiménez-Pastor and V. Pillwein, *A computable extension for D-finite functions: DD-finite functions* Bull. Soc. Math. France **47**, (1919), pp. 161-271.
- [49] A. Jiménez-Pastor, *DD-finite functions in Sage*, Séminaire Lotharingien de Combinatoire **82B** (2019) Article 101, 8pp.
- [50] P. Candelas, X. de la Ossa, P. Green and L. Parkes, *A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory*, Nucl. Phys. **B359**, (1991), pp. 21-74.
- [51] C. F. Doran, *Picard-Fuchs Uniformization and Modularity of the Mirror Maps*, Comm. Math. Phys. **212**, pp. 625-647, (2000).
- [52] C. F. Doran, *Picard-Fuchs Uniformization: Modularity of the Mirror Map and Mirror-Moonshine*, CRM Proc. Lecture Notes, **24**, Amer. Math. Soc. pp. 257-281, Providence and arXiv:math/9812162v1, (1998).
- [53] B.H. Lian and S-T. Yau, *Mirror Maps, Modular Relations and Hypergeometric Series II*, Nuclear Phys. **B 46**, Proceedings Suppl. Issues 1-3, (1996) pp.248-262 and arXiv: hep-th/950753v1 (1995)
- [54] C. Krattenthaler and T. Rivoal, *On the Integrality of the Taylor Coefficients of Mirror Maps*, Communications in Number Theory (2009), Volume: **3**, <http://www-fourier.ujf-grenoble.fr/~rivoal> and arXiv:0709.1432v3[math.NT]
- [55] S Boukraa and J-M Maillard, *Selected non-holonomic functions in lattice statistical mechanics and enumerative combinatorics*, 2016, J. Phys. **A 49**: Math. Theor (29 pages) 074001 and arXiv:1510.04651v1 [math-ph]
- [56] A. J. Guttmann, I. Jensen, J-M. Maillard, J. Pantone, *Is the full susceptibility of the square-lattice Ising model a differentially algebraic function ?*, (2016) J. Phys. **A 49**: Math. Theor. 504002 (36 pages) Special Issue in honour of A. J. Guttmann, and arXiv:1607.04168v2 [math-ph]
- [57] J. McKay and A. Sebbar, *Replicable functions: an introduction*, Springer
- [58] A. El Basraoui and J. McKay, *The Schwarzian equation for completely replicable functions*, LMS J. Comput. Math. **20** (1) (2017) 30-52
- [59] B. Heim and A. Murasea, *Completely replicable functions and symmetries*, Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg, (2019).
- [60] D. Ford, J. McKay and S. Norton, *More on replcable functions*, Communications in Algebra **22**, (1994) pp. 5175-5193 (2007)
- [61] A. El Basraoui, *Modular functions and replicable functions*, (2004)
- [62] D. Alexander, C. Cummins, J. McKay and C. Simons, *Completely replicable functions*, Groups, combinatorics and geometry (Durham, 1990), London Mathematical Society Lecture Note Series **165** (Cambridge University Press, Cambridge, 1992) 87-98.
- [63] B Heim, *Completely replicable functions and symmetries* Volume 89, pages 169-177, (2019) Springer

- [64] Chang Heon Kim and Ja Kyung Koo, *Super-replicable functions $N(j1, N)$ and periodically vanishing property*, J. Korean Math. Soc. 2007; 44(2): 343-371
- [65] G. Casale, *Enveloppe Galoisienne d'une application rationnelle de \mathbb{P}_1* , Publicacions Matemàtiques, Vol. **50**, No. 1 (2006), pp. 191-202 Published by: Universitat Autònoma de Barcelona, arXiv[math/0503424]
- [66] G. Casale, *An introduction to Malgrange pseudogroup*, SMF - Séminaires et Congrès **23** (2011)
- [67] G. Casale, *El grupoide de Galois de una transformación racional*, VIII Escuela Doctoral intercontinental de Matemáticas PUCP-UVa 2015 CIMPA Research school "Transformation Groups and Dynamical Systems"
- [68] G. Casale and Julien Roques, *Dynamic of rational mappings and difference Galois theory*, Int. Math. Res. Notices 2008 (2008)
- [69] G. Casale *Sur le groupoïde de Galois d'un feuilletage*, Thèse de doctorat effectuée sous la direction d'Emmanuel Paul et Jean-Pierre Ramis, soutenue le 09/07/2004
- [70] G. Casale, *D-enveloppe d'un difféomorphisme de $(C, 0)$* , Annales de la Faculté des Sciences de Toulouse, Mathématiques, Tome XIII, (2004) pp. 515-538.
- [71] G. Casale, *Morales-Ramis Theorems via Malgrange pseudogroup*, Tome 59, (2009), pp. 2593-2610., Annales de l'institut Fourier,
- [72] B. Malgrange, *On nonlinear differential Galois Theory*, Ann. of Math. 23B:2 (2002), pp. 219-226.
- [73] A. Bostan, S. Boukraa, S. Hassani, J.-M. Maillard, J.-A. Weil, N. Zenine, and N. Abarenkova, *Renormalization, isogenies and rational symmetries and differential equations*, Advances in Mathematical Physics, Volume **2010** (2010), Article ID 941560, 44 pages, <https://www.hindawi.com/journals/amp/2010/941560/>
- [74] A. Buium, *Geometry of differential polynomial functions III: moduli spaces*, Amer. J. Math. 117 (1995) 1-73.
- [75] Y. Abdelaziz, S. Boukraa, C. Koutschan and J.-M. Maillard, *Heun functions and diagonals of rational functions*, (2020) J.Phys. **A 53**: Math. Theor. 075206 (24p).
- [76] G. Shimura, *Correspondances modulaires et les fonctions ζ de courbes algébriques*, Journal of the Math. Soc. of Japan, **10**, (1958), pp. 1-27.
- [77] Y. Abdelaziz, S. Boukraa, C. Koutschan and J.-M. Maillard, *Heun functions and diagonals of rational functions (unabridged version)*, arXiv:1910.10761 [math-ph] (2019) (64p).
- [78] I. Bouw and M. Möller, *Differential equations associated with nonarithmetic Fuchsian groups*, arXiv:0710.5277v1 [math.AG] 2007
- [79] M. Dettweiler and S. Reiter, *On globally nilpotent differential equations*, Journal of Differential Equations **248**, 2010, pp. 2736-2745
- [80] B. Dwork, *Differential operators with nilpotent p -curvature*, American Journal of Mathematics **112** 1990 pp. 749-786
- [81] C. Krattenthaler and T. Rivoal, *Analytic properties of of Mirror Maps*, J. Aust. Math. Soc. **92** (2012) pp.195-235
- [82] Y. Abdelaziz, C. Koutschan and J.-M. Maillard, *On Christol's conjecture*, 2020 J. Phys. A: Math. Theor. 53 205201
- [83] A. Bostan, S. Boukraa, S. Hassani, J.-M. Maillard, J.-A. Weil, and N. Zenine, *Globally nilpotent differential operators and the square Ising model*, J. Phys. A: Math. Theor. **42** (2009) 125206 (50pp) and arXiv:0812.4931
- [84] S. Boukraa et J.-M. Maillard, *The lambda extension of the Ising correlation functions $C(M, N)$* , 2023, J. Phys. A: Math. Theor. **56** 085201 and arXiv:2209.07434v2
- [85] S. Boukraa, C. Cosgrove, J.-M. Maillard, and B. M. McCoy, *Factorization of Ising correlations $C(M, N)$ for $\nu = -k$ and $M + N$ odd, $M \leq N$, $T < T_c$ and their lambda extensions* 2022 J. Phys. A: Math. Theor. **55** 405204
- [86] M. van Hoeij, R. Vidunas, *Belyi functions for hyperbolic hypergeometric-to-Heun transformations*, Journal of Algebra (2015), Volume: **441**, pp. 609-659, arXiv:1212.3803v3[math.AG]
- [87] J. Sijsling and J. Voight, *On computing Belyi maps*, Publications mathématiques de Besançon (2014), Issue: 1, pp.73-171, arXiv:1311.2529v3[math.NT]
- [88] R. Vidunas, A. V. Kitaev, *Computation of highly ramified coverings*, Math. Comp. (2009), Volume: **78**, pp.2371-2395, arXiv:0705.3134[math.AG]
- [89] L. S. Khadjavi and V. Scharaschkin, *Belyi maps and Elliptic Curves*, <http://myweb.lmu.edu/lkhadjavi/belyielliptic.pdf>
- [90] D. Masoero, *Painlevé I, Coverings of the Sphere and Belyi Functions*, Constr Approx (2014), Volume: **39**, Issue: 1, pp.43-74, arXiv:1207.4361v2[math.ph]
- [91] R. Maier, *On reducing the Heun equation to the hypergeometric equation*, J. Differential Equations **213** (2005), no. 1, pp. 171-203.
- [92] A.J. Guttmann and J.-M. Maillard, *Automata and the susceptibility of the square lattice Ising*

- model modulo powers of primes*, (2015) J. Phys. **A 48**: Math. Theor (22 pages) 47001, Special Issue dedicated to R.J. Baxter and arXiv:1507.02872v2 [math-ph]
- [93] S. Boukraa, A.J. Guttmann, S. Hassani, I. Jensen, J.-M. Maillard, B. Nickel and N. Zenine, *Experimental mathematics on the magnetic susceptibility of the square lattice Ising model*, J. Phys. A: Math. Theor. **41** (2008) 455202 (51pp) and arXiv:0808.0763